# Comparison of one repetition maximum performance across three weightlifting overhead pressing exercises and sport groups

# 4 ABSTRACT

Objective: This study aimed to (I) compare the one repetition maximum (1RM) 5 6 performance between the push press (PP), push jerk (PJ) and split jerk (SJ), and (II) explore these differences between weightlifters, CrossFit<sup>®</sup> athletes and a mixed group of 7 athletes. *Method:* Forty-six resistance trained male (age:  $28.8 \pm 6.4$  years; height: 180.0 8 9  $\pm$  6.0 cm; body mass: 84.1  $\pm$  10.2 kg; weightlifting training experience: 3.64  $\pm$  3.14 years) 10 participated in this study. The 1RM performance of the PP, PJ and SJ were assessed during the same session in a sequential order (i.e. combined 1RM assessment method). 11 Thirty-six participants were re-tested to determine between-session reliability of the 1RM 12 13 values. Results: Intraclass correlation coefficients (ICC) and associated 95% confidence intervals showed a high between-session reliability for the PP (ICC = 0.98 [0.95-0.99]), 14 15 PJ (ICC = 0.99 [0.98-1.00]) and SJ (ICC = 0.99 [0.98-1.00]). There was a significant main effect of exercise ( $\eta^2 = 0.662$ ) and exercise x group interaction ( $\eta^2 = 0.066$ ) on the 1RM 16 performance (p<0.0001), while the main effect of group did not reach statistical 17 significance (p=0.072). *Conclusion:* This study provides evidence that the weightlifting 18 overhead pressing derivatives impact the 1RM performance. In addition, the interaction 19 of exercise and sport group was caused by the higher differences in the 1RM performance 20 21 between-exercises for weightlifters compared to CrossFit<sup>®</sup> and a mixed group of athletes. 22 Therefore, strength and conditioning professionals should be aware that the differences 23 in 1RM performance between weightlifting overhead pressing derivatives may be 24 affected by the sport group.

25

26 Key words: push press, push jerk, split jerk, maximal strength, combined assessment

#### 27 INTRODUCTION

High levels of muscular strength and power development are key to maximizing athletic
performance.<sup>1</sup> Researchers currently suggest that greater muscular strength underpins
many physical and performance attributes and can be vastly influential in the ability to
generate high rates of force development (RFD) and high levels of power.<sup>1,2</sup> Furthermore,
James et al.<sup>3</sup> found that improvements in performance of high velocity sporting actions
are associated with pre-existing strength levels.

34

35 The one repetition maximum (1RM) is considered the gold standard for assessing the maximal strength during dynamic tasks.<sup>4</sup> The 1RM can be performed using the same 36 exercises as those undertaken by the athletes during their regular training and is 37 38 commonly used to prescribe the loads during resistance training programs.<sup>5</sup> The 1RM is frequently assessed by strength and conditioning professionals to compare the strength 39 40 between sports and to monitor the changes in strength over time; however, only a few exercises such as the bench press, squat and power clean are commonly reported.<sup>6,7</sup> For 41 example, McBride et al.<sup>7</sup> found that the 1RM performance in the squat exercise was able 42 43 to discriminate the strength levels of weightlifters, powerlifters, sprinters and 44 recreationally trained athletes (p < 0.05). Of note is that the weightlifter's 1RM was 45 significantly higher compared to sprinters 1RM (p < 0.05) but no significant differences 46 were observed compared to powerlifters (p>0.05).

47

48 The fact that the bench press, squat and power clean are the exercises most commonly 49 used for assessing strength levels may lead to insufficient and inaccurate information to provide strength guidelines to various populations across numerous exercises included in 50 51 resistance training programs. It has been suggested that multiple measurements of strength should be used to capture general strength adaptations of a given athlete.<sup>4</sup> It is 52 53 important to note that there are notable differences in 1RM performances between 54 exercises with similar kinematics. For example, Kelly et al.<sup>8</sup> compared the 1RM 55 performance between three power clean variations (from the floor, from the knee and 56 from mid-thigh) in twelve male collegiate athletes, reporting subtleties in exercise 57 technique resulting in meaningful differences (>6%) in the 1RM performance. Similarly, 58 Yavuz et al.<sup>9</sup> reported meaningful differences between the front and back squat variations 59 with participants lifting significantly higher loads during the back squat compared to the 60 front squat (p < 0.05).

61

62 Researchers have suggested potential benefits for implementing overhead pressing 63 derivatives as training tools to improve not only weightlifting performance, but also to enhance sports performance.<sup>10</sup> Notably, researchers have provided evidence that these 64 exercises may be a time-effective method of enhancing lower-body strength and power; 65 66 however, modifications in technical execution and equipment result in meaningful differences in force-time characteristics.<sup>11,12</sup> Therefore, the assessment of 1RM 67 performance of different overhead pressing variations may provide valuable information 68 69 to practitioners. To the authors' knowledge, there is little information regarding 1RM 70 overhead pressing derivatives and no research has compared differences in 1RM between 71 the main overhead pressing derivatives as the push press (PP), push jerk (PJ), and split 72 jerk (SJ). Furthermore, it is important to note that although researchers have compared the 1RM performance between variations of the power clean,<sup>8</sup> they have not compared 73 74 these exercise variations between athletes of different sporting backgrounds.

75

The main aim of this study was to compare the 1RM performance between three 76 variations of overhead pressing (PP, PJ and SJ) for three different sport populations 77 (weightlifters, CrossFit<sup>®</sup> athletes and a mixed group of athletes). A further aim of the 78 study was to compare to 1RM performance of these variations across the three sport 79 populations. Based on the evidence that the greatest force, velocity and power occur 80 81 during the SJ,<sup>10</sup> it was hypothesized that, regardless of the sport population, the 1RM 82 performance would be ranked from the highest to the lowest as follows: SJ, PJ and PP variation. Due to the higher predominance of overhead pressing derivatives in 83 weightlifters' training routines,<sup>13</sup> it was also hypothesized that weightlifters would 84 85 present the highest 1RM for these exercises.

86

## 87 METHOD

88 Subjects

89 Forty-six resistance trained males volunteered to participate in this study (Table 1). 90 Participants were amateur competitors in regional and national tournaments in their 91 respective disciplines. Furthermore, they were required to have at least six months of weightlifting experience including the PP, PJ and SJ, performed regularly ( $\geq$  3 times per 92 93 week) in their respective strength and conditioning training preparations. Participants 94 were subdivided into three groups depending on their sport background: weightlifting, CrossFit<sup>®</sup>, and a mixed group of athletes (e.g. rugby league, rugby union, football, track 95 and field, wrestling, gymnastic, volleyball and basketball). All participants had 96 97 previously performed 1RM testing for a variety of exercises. The investigation was 98 approved by the institutional review board of the University of Salford, and all 99 participants provided written informed consent before participation. The study conformed 100 to the principles of World Medical Association's Declaration of Helsinki. Participants were supervised by a certified strength and conditioning specialist during all testing 101 102 sessions to allow familiarization with the protocols and ensure appropriate technique.

103

## 104 Experimental design

105 A within- and between-subjects design was adopted to determine the between session 106 reliability of 1RM performance and to determine the differences in the 1RM performance 107 across three weightlifting overhead pressing derivatives (PP, PJ and SJ), while also comparing across sport groups (weightlifters, CrossFit<sup>®</sup> and a mixed group of athletes). 108 46 healthy men were evaluated using the combined 1RM assessment method; a 109 110 standardized protocol previously validated on a similar population for overhead pressing exercises <sup>14</sup>. In brief, the 1RM performance of the PP, PJ and SJ were assessed during the 111 112 same session in a sequential order. All testing sessions were performed using standardized barbells and plates (Werksan weights and Olympic bar; Werksan, Moorestown, New 113 Jersey, USA), lifting platforms and power racks (Powerlift, Iowa, USA). Furthermore, 114 115 thirty-six participants performed the combined 1RM assessment method one week apart, 116 to test the between-session reliability of the 1RM performance for each exercise. Verbal encouragement was provided throughout all maximal testing conditions. Participants 117 were asked to replicate their fluid and food intake twenty-four hours before each day of 118 119 testing, to avoid strenuous exercise for forty-eight hours before testing, and to maintain any existing supplementation regimen throughout the duration of the study. All testing 120 was performed at the same time of day for all participants to minimize the effect of 121 122 circadian rhythms.

123

124 *Testing procedures* 

Participants completed a warm up protocol which has been previously described by Soriano et al.<sup>14</sup> Briefly, this consisted of dynamic activation, exercise-specific drills and one set of five submaximal (50-60% of the maximal perceived effort) repetitions in each exercise (PP, PJ and SJ). Five minutes of rest was taken, followed by another set of three submaximal (70-85% of the maximal perceived effort) repetitions for each exercise. After the warm-up, participants rested for 5 min before the start of the combined 1RM assessment method.

- 132
- 133 Measurement equipment and data analysis.

134 The combined 1RM assessment consisted of performing the 1RM test for the PP, PJ and SJ in a sequential order. The three exercises started from a near-maximal load (95% of 135 the maximal perceived effort) and each successful attempt was followed by a small 136 increment of the load (2.5-5.0%) until the 1RM was reached, following previous NSCA 137 138 guidelines.<sup>15</sup> Participants rested for 3 to 5 min between attempts within the same exercise 139 and between-exercises. Hence, the 1RM in PP served as a preparation exercise for the PJ 140 and both for the SJ, due to the fact that all of these exercises have a similar movement pattern.<sup>10</sup> The barbell was taken out of power racks before starting each attempt to 141 142 minimize the fatigue associated with the performance of the clean, which precedes the jerk in competitions.<sup>16</sup> 143

144

## 145 *Statistical analyses*

Descriptive data are presented as means and standard deviation (SD). To determine the 146 between-session reliability of the 1RM assessment, the coefficient of variation (CV), 147 148 intraclass correlation coefficient (ICC; model 3.1) and associated 95% confidence intervals (CI) were calculated and interpreted based on the recommendations of Cortina<sup>17</sup> 149 150 where an ICC > 0.80 is considered as highly reliable, while a CV < 15% was used as a criterion for the minimum acceptable reliability.<sup>18</sup> The reliability analysis was performed 151 by means of a custom spreadsheet,<sup>19</sup> while SPSS (version 25.0: SPSS, Inc., Chicago, IL, 152 USA) was used for the remaining analyses. Normality of the anthropometric measures 153 154 (height and body mass) for each group was confirmed by the Shapiro-Wilk's (p>0.05), 155 while the age and weightlifting training experience were not normally distributed for weightlifters and mixed athlete groups (p<0.05). Two one-way analysis of variance 156 (ANOVA) with Bonferroni post hoc analysis were used to test between-group differences 157 158 for the height and body mass, while the Kruskal-Wallis' test for independent samples was 159 used to compare the age and weightlifting experience between groups.

160

Shapiro-Wilk's and Levene's tests were used to determine the distribution of the 1RM 161 performances and the homogeneity of variances, respectively. The Greenhouse-Geisser 162 163 correction was used when the assumption of sphericity was violated (p < 0.05). A mixed repeated measures ANOVA with Bonferroni post hoc analysis was applied using the 164 exercises (PP, PJ, and SJ) as within-subject factor, and group (weightlifters, CrossFit<sup>®</sup>, 165 and mixed athlete groups) as between-subject factor. An *a priori* alpha level was set at p 166 167 < 0.05. Effect sizes were determined using eta squared ( $\eta^2$ ) to determine the magnitude of the effect independently of the sample size;  $\eta^2$  has previously been recommended for 168 ANOVA designs,<sup>20</sup> and interpreted based on the recommendations of Cohen,<sup>21</sup> where 169 effect sizes of < 0.06 are considered small, 0.06-0.14 are medium, and > 0.14 are 170 171 considered large effects.

172

#### 173 **RESULTS**

The ICCs demonstrated high between-session reliability of the combined 1RM assessment for the PP (ICC = 0.98 [0.95-0.99]), PJ (ICC = 0.99 [0.98-1.00]) and SJ (ICC = 0.99 [0.98-1.00]). In addition, the low CV also confirmed the high reliability for the PP (CV = 2.45 % [1.99-3.20]), PJ (CV = 1.57 % [1.27-2.05]) and SJ (CV = 1.67 % [1.35-2.17]).

179

Participants characteristics' mean and SD (age, height, body mass and weightlifting training experience) are shown in Table 1. The results of the one-way ANOVA revealed no significant differences for the anthropometric measures (height and body mass), while the Kruskal-Wallis test revealed significant (p=0.004) differences for the age betweengroups. The results of the Mann-Whitney U's test revealed that the CrossFit® group was significantly (p=0.004) older than the weightlifters and the mixed group of athletes. In addition, no differences were found for weightlifting training experience between groups.

188

Table 1.	Descriptive	characteristics	of the g	groups
----------	-------------	-----------------	----------	--------

Group	Sample size (n)	Age (years)	Height (cm)	Body mass (kg)	WL training experience (years)
WL	15	26.4 ± 1.6	177.9 ± 1.8	81.1 ± 3.1	4.5 ± 1.3
[range]		[from 19 to 40]	[from 166 to 193]	[from 62 to 108]	[from 0.5 to 20]
CF	19	32.1 ± 1.1*	180.4 ± 1.3	83.4 ± 2.0	3.3 ± 0.3
[range]		[from 22 to 42]	[from 169 to 191]	[from 68 to 99]	[from 1 to 7]
MX	12	26.2 ± 2.1	182.1 ± 1.2	88.9 ± 2.5	3.2 ± 0.6
[range]		[from 20 to 43]	[from 174 to 190]	[from 77 to 105]	[from 0.8 to 7.5]

WL = weightlifting, CF = CrossFit<sup>®</sup> athletes, MX = mixed group of athletes, \*significantly (p=0.004) higher than the WL and MX group

191 The results of the ANOVA revealed a significant main effect of exercise and exercise x 192 group interaction on the 1RM performance with medium and small effect sizes, 193 respectively. However, for the combined 1RM performances there were no significant 194 differences between groups with an observed power of 0.918 (Table 2).

195

Table 2. ANOVA's results of comparing within- and between-subjects' effects on the 1RM performance

Exercise -	Group			Main effect		Interaction	
	WL	CF	MX	Group	Exercise	Exercise x Group	
PP (kg)	94.7 ± 17.6	86.1 ± 13.8	90.4 ± 14.8	F = 1.8 p = 0.175	F = 105.6 p < 0.001	F = 6.0 p < 0.001	
PJ (kg)	105.5 ± 17.9	93.0 ± 14.0	97.1 ± 17.4				
SJ (kg)	115.3 ± 21.2	96.6 ± 16.3	102.9 ± 18.3	<b>ग</b> - = 0.000	η- = 0.101	η- = 0.012	

PP = push press, PJ = push jerk, SJ = split jerk, WL = weightlifters, CF = CrossFit<sup>®</sup> athletes, MX = mixed group of athletes, F = F value, p = significance value;  $\eta^2$  = eta squared

196 197

Weightlifters demonstrated significantly higher 1RM SJ performance  $(115.3 \pm 21.2 \text{ kg})$ compared to the PJ  $(105.5 \pm 17.9 \text{ kg}; \text{ p} < 0.001)$  and PP  $(94.7 \pm 17.6 \text{ kg}; \text{ p} < 0.001)$ . The

<sup>189</sup> 190

CrossFit® group demonstrated significantly higher 1RM SJ performance  $(96.6 \pm 16.3 \text{ kg})$ compared to the PP ( $86.1 \pm 13.8 \text{ kg}$ ; p<0.001), but not significant than the PJ ( $93 \pm 14 \text{ kg}$ ; p=0.90). The mixed group of athletes demonstrated significantly higher 1RM SJ performance ( $102.9 \pm 18.3 \text{ kg}$ ) compared to the PP ( $90.4 \pm 14.8 \text{ kg}$ ; p<0.001), but not significantly greater than the PJ ( $97.1 \pm 17.4 \text{ kg}$ ; p=0.110) (Fig 1).

205



206

Figure 1. Comparison of one repetition maximum performances between exercises and between groups, PP
push press, PJ push jerk, SJ split jerk, WL weightlifters, CF CrossFit® athletes, MX mixed group of
athletes, a significantly (p<0.001) lower than the WL-SJ, b significantly (p<0.001) greater than the CF-PP,</li>
c significantly (p<0.001) greater than the MX-PP, \*significantly (p<0.001) greater than the WL-PP, #</li>
Significantly (p<0.001) greater than CF-PP, \$significantly (p=0.028) greater than MX-PP.</li>

212

#### 213 **DISCUSSION**

The main finding of this study was that a significant effect of exercise was found on the IRM performance in line with our previous hypothesis, however, there was no significant effect of group. These findings are important for strength and conditioning coaches because they describe not only the variations of the 1RM performance through the main overhead pressing derivatives (PP, PJ and SJ), but also how these variations occurred in different sport groups. In addition the ICC and CV showed a high between-session reliability for the three exercises in line with previous results.<sup>14</sup>

221

As hypothesized, the SJ was the exercise where the heaviest loads were lifted for the three 222 groups, followed by the PJ and PP. However, for the CrossFit<sup>®</sup> athletes and the mixed 223 group there were no significant differences for the 1RM performance between the PJ and 224 SJ. This 1RM performance sequence was expected based on the notion that the jerk 225 provides a greater stimulus for force, velocity and power production.<sup>10,22</sup> The differences 226 in the 1RM performance may be attributable to the technical peculiarities of each exercise 227 variation as previously stablished for the power clean.<sup>8</sup> According to this, although the 228 229 three exercises share the same lower-body propulsion pattern (composed by the dip, breaking and thrusting phases), there are important differences between the PP, PJ and SJ 230 at the end of the propulsion phase. Mainly, in the PP the lifter must continue pressing the 231 barbell with the upper body (flexion of the shoulders and extension of the elbows) to 232 reach the overhead position in a continuous motion;<sup>23</sup> on the other hand, when the barbell 233 leaves the shoulders during the PJ and SJ the lifter must drop under the barbell in either 234 a <sup>1</sup>/<sub>4</sub> squat (PJ) or split position (SJ) to catch the barbell overhead.<sup>24</sup> Therefore, the SJ and 235

PJ are clearly advantageous to lift greater loads as less barbell displacement and therefore
 mechanical work is required to receive the barbell overhead.<sup>10,24</sup>

237 238

The groups included in this study represent 3 different sport populations based on their 239 240 competitive disciplines. Previously, researchers found that maximal strength levels of the 241 lower body extremities may discriminate the long-term adaptations of the training specificity in athletes from different sports.<sup>7,25</sup> For example, Izquierdo et al.<sup>25</sup> found that 242 a weightlifting group was differentiated in comparison to other sport groups (e.g. handball 243 players, middle-distance runners, road cyclists and a control group), by their higher 1RM 244 245 performance in the half squat; however, McBride et al.<sup>7</sup> found that 1RM performance was able to differentiate weightlifters vs. sprinters and a control group, but not from the 246 247 powerlifters. Since the training routines of both weightlifters and powerlifters usually 248 implement heavy squats, it seems that the training specificity plays a major role of the 249 training adaptations and therefore, the 1RM performance of a given exercise is influenced by such training background. In the present study, all participants had weightlifting 250 251 training experience and no statistical differences in weightlifting training experience were 252 found between groups. Although there were no significant differences in the 1RM 253 performance between groups, the weightlifters showed a trend to have higher 1RM 254 performances for the three exercises in comparison to the other groups. However, our results show that the PP, PJ and SJ were not able to discriminate between groups as 255 previously stated for the squat.<sup>7,25</sup> Note that differences in training experience were based 256 on vears of training, however the hours of training and practice in the weightlifters are 257 258 likely to be substantially higher than in other groups.

259

260 A clear interaction effect between exercises and sport group was found in the present 261 study. For example, the weightlifters' SJ was the most discriminative exercise as this 262 presented the biggest difference in performance between the three groups; however, no significant differences were found between the PJ and SJ for the CrossFit<sup>®</sup> and the mixed 263 264 group of athletes. Although these differences may be attributable to a wide range of physiological and psychological factors,<sup>16,26,27</sup> these results support the specificity 265 training principle<sup>7,25</sup> and the theory of Buckner et al.<sup>4</sup>. The theory of Buckner et al.<sup>4</sup> 266 reflects that the 1RM performance assessment (test) is a highly specific sporting task, 267 which means that the more the practice, the better the performance.<sup>4</sup> Due to that the SJ is 268 highly trained by weightlifters and is part of official weightlifting competitions,<sup>13,16</sup> its 269 270 discriminative role only for the weightlifters group is justified.

271

Finally, the training status may be a limiting factor in the generalizability of these findings.<sup>4,28</sup> Note that although the three groups can be considered as well-trained athletes due to the specified resistance training background of  $\geq 2$  years,<sup>29</sup> they may otherwise be classified as recreationally trained based on the recommendations of Rhea.<sup>30</sup> Therefore, more research is needed to elucidate how world class and more experienced participants than those used for this study may impact the 1RM performance during the PP, PJ and SJ.

- 279
- 280 281

## 282 PRACTICAL APPLICATION

283 The present study provides evidence that the main weightlifting overhead pressing 284 derivatives impact the 1RM performance. Strength and conditioning professionals should 285 be aware of these differences in order to prescribe the loads adequately and choose the

desired exercises. The SJ may be more favourable when attempting to increase the 286 maximal strength levels in weightlifters, followed by PJ and PP. However, CrossFit<sup>®</sup> and 287 the mixed group of athletes may use the SJ and PJ interchangeably when the objective is 288 to increase the maximal strength levels. Furthermore, this research extends the specificity 289 principle due to the interaction effect of exercise (PP, PJ and SJ) and sport group on the 290 291 1RM performance. Specifically, weightlifters presented higher differences in the 1RM performance between-exercises (9-22%), in comparison to the CrossFit<sup>®</sup> group (4-11%) 292 and the mixed group of athletes (6-14%). Therefore, practitioners should be aware of 293 differences in the 1RM performance between weightlifting overhead pressing derivatives 294 295 may be affected by the sport group. This study provides preliminary evidence for future research into a comparison of the variations in the 1RM performances in a long-term 296 297 structured programme, where the training specificity of each group may have a greater 298 impact.

299 300

## 301 CONCLUSIONS

There is an effect of exercise on the 1RM performance during the main weightlifting overhead pressing derivatives (PP, PJ and SJ). The SJ is the exercise where the greatest loads were lifted for the three sport groups assessed; however, it was not significantly greater for the CF and MX groups in comparison to the PJ. In addition, the interaction of exercise and sport group revealed that the sport group impacted the 1RM performance with weightlifters showing the largest differences in 1RM performance between exercises.

- 309
- 310
- 311 312

## 313 ACKNOWLEDGMENTS

The authors thank the individuals, clubs and institutions (University of Salford, House of Lifters, CrossFit Salford, CrossFit Consilium, Force Athletes, Team Manchester Weightlifting, NorthWest Box, CrossFit Uros and CrossFit Bellum) which participated in this investigation. No grant funding was received to support this research, and the authors declare no conflict of interest.

- 319
- 320
- 321
- 322 323
- 323 324
- 325
- 326

# 327 **REFERENCES**

- Suchomel TJ, Nimphius S, Stone MH. The importance of muscular strength in athletic performance. *Sport Med.* 2016;46:1419-1449.
- Stone MH, Moir G, Glaister M, Sanders R. How much strength is necessary? *Phys Ther Sport*. 2002;3:88-96.
- 3. James LP, Haff GG, Kelly VG, Connick MJ, Hoffman BW, Beckman EM. The
  impact of strength level on adaptations to combined weightlifting, plyometric, and
  ballistic training. *Scand J Med Sci Sport*. 2018;28:1494-1505.
- 335 4. Buckner SL, Jessee MB, Mattocks KT, et al. Determining strength: a case for

336		multiple methods of measurement. Sport Med. 2017;47:193-195.
337	5.	Banyard HG, Nosaka K, Haff GG. Reliability and validity of the load-velocity
338		relationship to predict the 1RM back squat. J Strength Cond Res. 2017;31:1897-
339		1904.
340	6.	McMaster DT, Gill N, Cronin J, McGuigan M. A brief review of strength and
341		ballistic assessment methodologies in sport. Sport Med. 2014;44:603-623.
342	7.	Mcbride JM, Triplett-Mcbride T, Davie A, Newton RU, A comparison of strength
343		and power characteristics between powerlifters, Olympic lifters, and sprinters. J
344		Strength Cond Res. 1999;13:58-66.
345	8.	Kelly J, J. McMahon J, Comfort P. A comparison of maximal power clean
346		performances performed from the floor, knee and mid-thigh. J Trainology.
347		2015:3:53-56.
348	9.	Yavuz HU, Erdağ D, Amca AM, Aritan S. Kinematic and EMG activities during
349		front and back squat variations in maximum loads. J Sports Sci. 2015:33:1058-
350		1066.
351	10.	Soriano MA. Suchomel T. Comfort P. Weightlifting overhead pressing
352	101	derivatives: a review of the literature. <i>Sport Med.</i> 2019:49:867-885.
353	11.	Winwood PW, Cronin JB, Brown SR, Keogh J. A biomechanical analysis of the
354		strongman log lift and comparison with weightlifting's clean and jerk. Int J Sports
355		Sci Coach 2015:10:869-886
356	12	Renals I. Lake I. Keogh I. Austin K. Strongman log nush press: the effect log
357	12.	diameter has on force-time characteristics <u>J Strength Cond Res</u> 2018:32:2693-
358		2700
359	13	Stone MH Pierce KC Sands WA Stone MF Weightlifting: program design
360	15.	Strength Cond I 2006:28:10-17
361	14	Soriano MA García-Ramos A Torres-González A Castillo-Palencia I Marín PI
362	1	Comfort P. Validity and reliability of a standardized protocol for assessing the one
363		repetition maximum performance during overhead pressing exercises <i>I Strength</i>
364		Cond Res 2019 Epub ahead of print
365	15	Baechle TR Earle RW Essentials of strength training and conditioning (3 <sup>rd</sup>
366	10.	edition) Human kinetics 2008
367	16	Storey A Smith HK Unique aspects of competitive weightlifting: performance
368	10.	training and physiology Sport Med 2012:42:769-790
369	17	Corting IM What is coefficient alpha? An examination of theory and applications
370	17.	I Appl Psychol 1993.78.98
371	18	Baumgartner TA Chung H Confidence limits for intraclass reliability
372	10.	coefficients Meas Phys Educ Exerc Sci 2001:5:179-188
372	19	Hopkins W Calculations for reliability (Excel spreedsheet) A new view of
373	17.	statistics 2000
374	20	Lakens D. Calculating and reporting effect sizes to facilitate cumulative science: a
376	20.	practical primer for t-tests and ANOVAs <i>Front Psychol</i> 2013:4:863
370	21	Cohen I Statistical power analysis for the behavioral science (2 <sup>nd</sup> edition)
378	21.	Routledge 1988
370	22	Garhammer I. A comparison of maximal power outputs between elite male and
380	<i>LL</i> .	female weight lifters in competition Int I Snort Riomach 1991.7.3-11
281	23	Bishon C Chavda S Turner & Exercise technique: the push press Strangth Cond
282	23.	I 2018.40.104_108
202	24	Waller M Piner T Miller I Overhead pressing power/strength movements
207	∠+.	Strength Cond I 2000-21-20 40
304	25	Sirenzin COIR J. 2007, J1. J7 - 77.

385 25. Izquierdo M, Häkkinen K, Gonzalez-Badillo JJ, Ibáñez J, Gorostiaga EM. Effects

- of long-term training specificity on maximal strength and power of the upper and
  lower extremities in athletes from different sports. *Eur J Appl Physiol*.
  2002;87:264-271.
- 389 26. Abernethy P, Wilson G, Logan P. Strength and power assessment: issues, controversies and challenges. *Sport Med.* 1995;20:205.
- 391 27. Suchomel TJ, Nimphius S, Bellon CR, Stone MH. The importance of muscular
  392 strength: training considerations. *Sport Med.* 2018;765-785.
- Ritti-Dias RM, Avelar A, Salvador EP, Cyrino ES. Influence of previous
  experience on resistance training on reliability of one-repetition maximum test. J *Strength Cond Res.* 2011;25:1418-1422.
- 29. Cormie P, McGuigan MR, Newton RU. Influence of strength on magnitude and
  mechanisms of adaptation to power training. *Med Sci Sports Exerc*. 2010;42:15661581.
- 30. Rhea MR. Determining the magnitude of treatment effects in strength training
  research through the use of the effect size. *J Strength Cond Res*. 2004;18:918-920.
- 401
- 402 403

## 404 FIGURE CAPTIONS

Figure 1. Comparison of one repetition maximum performances between exercises and between groups, PP
push press, PJ push jerk, SJ split jerk, WL weightlifters, CF CrossFit® athletes, MX mixed group of
athletes, a significantly (p<0.001) lower than the WL-SJ, b significantly (p<0.001) greater than the CF-PP,</li>
c significantly (p<0.001) greater than the MX-PP, \*significantly (p<0.001) greater than the WL-PP, #</li>
Significantly (p<0.001) greater than CF-PP, \$significantly (p=0.028) greater than MX-PP.</li>