

DNA metabarcoding unveils multi-scale trophic variation in a widespread coastal opportunist

Andjin Siegenthaler^{1,3,*}, Owen S. Wangensteen^{1,4}, Chiara Benvenuto¹, Joana Campos² & Stefano Mariani¹

¹ Ecosystems and Environment Research Centre, School of Environment & Life Sciences University of Salford, UK

² CIMAR/CIIMAR - Centro Inter disciplinar de Investigação Marinha e Ambiental, Universidade do Porto, Portugal

³ Current address: Department of Integrative Biology, University of Guelph, Guelph, Canada

⁴ Current address: Norwegian College of Fishery Science, UiT the Arctic University of Norway, Tromsø, Norway

* Corresponding author:
Andjin Siegenthaler
School of Environment and Life Sciences
Room 335, Peel Building, University of Salford,
Salford M5 4WT, UK
t: +44(0)1612953136
e: a.siegenthaler@edu.salford.ac.uk

Running title: Brown shrimp diet metabarcoding

Keywords: Community Ecology; Crustaceans; Diet Analysis; Metabarcoding; Foodwebs; Environmental DNA

Abstract

A thorough understanding of ecological networks relies on comprehensive information on trophic relationships among species. Since unpicking the diet of many organisms is unattainable using traditional morphology-based approaches, the application of high-throughput sequencing methods represents a rapid and powerful way forward. Here, we assessed the application of DNA-metabarcoding with nearly universal primers for the mitochondrial marker cytochrome c oxidase I (COI) in defining the trophic ecology of adult brown shrimp, *Crangon crangon*, in six European estuaries. The exact trophic role of this abundant and widespread coastal benthic species is somewhat controversial, while information on geographical variation remains scant. Results revealed a highly opportunistic behaviour. Shrimp stomach contents contained hundreds of taxa (>1000 molecular operational taxonomic units), of which 291 were identified as distinct species, belonging to 35 phyla. Only twenty ascertained species had a mean relative abundance of more than 0.5 %. Predominant species included other abundant coastal and estuarine taxa, including the shore crab *Carcinus maenas* and the amphipod *Corophium volutator*. Jacobs' selectivity index estimates based on DNA extracted from both shrimp stomachs and sediment samples were used to assess the shrimp's trophic niche indicating a generalist diet, dominated by crustaceans, polychaetes and fish. Spatial variation in diet composition, at regional and local scales, confirmed the highly flexible nature of this trophic opportunist. Furthermore, the detection of a prevalent, possibly endoparasitic fungus (*Purpureocillium lilacinum*) in the shrimp's stomach demonstrates the wide range of questions that can be addressed using metabarcoding, towards a more robust reconstruction of ecological networks.

Introduction

Trophic interactions provide important insights on a wide range of ecological dynamics, ranging from individual to ecosystem levels, which include animal behaviour, predator-prey interactions, food web structure and community ecology (e.g. Leray *et al.* 2015; Pinol *et al.* 2014). The feeding strategy of key consumers can have pronounced influences on ecosystem dynamics (Hanski *et al.* 1991; Holling 1965) and their stomach contents can reveal essential information on food item distribution and prey assemblage structure (Lasley-Rasher *et al.* 2015). Crustaceans are a key component in marine/estuarine soft bottom habitats (Evans 1983, 1984) and evaluating their diet is very challenging due to the complexity of direct observations on predation rates and the limitations associated with the identification of partially digested food items (Asahida *et al.* 1997; Feller 2006; Symondson 2002).

The recent application of high-throughput sequencing (HTS) tools, such as metabarcoding, promises to revolutionise the way prey diversity and composition are estimated from gut contents or faeces of consumers (Kartzinel & Pringle 2015; Leray *et al.* 2015). Metabarcoding refers to the identification of multiple taxa based on the screening of bulk DNA extracted from community or environmental samples (i.e. water, soil, faeces; Barnes & Turner 2016), by means of massive parallel sequencing of PCR amplicons (Barnes & Turner 2016; Taberlet *et al.* 2012a). Metabarcoding has proven to be highly effective for the identification of prey remains with improved taxonomic resolution, accuracy and speed of analysis, compared to traditional morphological methods (Berry *et al.* 2015; Casper *et al.* 2007; Symondson 2002). Yet, some challenges remain, such as fragmentation of partially-digested DNA, variability in taxon-specific digestion rates, secondary predation, and, typically, the presence of high proportion of DNA from the study organisms itself, which may reduce sequencing depth and render cannibalism undetectable (Barnes & Turner 2016; Berry *et al.* 2015; Pinol *et al.* 2014). Furthermore, due the sensitivity of these methods, in some cases, it might be difficult to discriminate between contaminant DNA and target DNA.

The brown shrimp, *Crangon crangon* (L.) is a key crustacean species in European coastal waters. Its wide distribution (i.e. from the White Sea to Morocco), year round occurrence, and high abundance ($> 100 \text{ ind. m}^{-2}$; van der Veer *et al.* 1998) make it an essential part of the coastal benthic food web (Ansell *et al.* 1999; Campos & van der Veer 2008; Evans 1984), a major prey item for birds and fish (Evans 1984; Walter & Becker 1997), and an important target for fisheries, with recorded catches in 2011 up to 35,000 tons and more than 500 fishing vessels employed in the North Sea (Aviat *et al.* 2011; Campos & van der Veer 2008). The trophic position of *C. crangon* is still being discussed, being described as trophic generalist (Evans 1983), carnivorous opportunist (Pihl & Rosenberg 1984) omnivorous (Ansell *et al.* 1999; Raffaelli *et al.* 1989; Tiews 1970) and probable scavenger (Ansell *et al.* 1999). As a juvenile, it relies mostly on the consumption of meiofaunal prey items while it switches to larger demersal organisms as an adult, including conspecifics and juvenile stages of several commercially important teleosts and bivalves (Evans 1984; Oh *et al.* 2001; Pihl & Rosenberg 1984; van der Veer & Bergman 1987; van der Veer *et al.* 1998). Previous studies showed considerable variation in prey item consumption, partly due to the brown shrimp's inherent trophic flexibility and nice breath, but also because studies have relied on microscopic identification of prey remains (e.g. Boddeke *et al.* 1986; Oh *et al.* 2001; but see also Nordström *et al.* 2009); yet, prey items are usually macerated to a fine degree by *C. crangon*, and a high proportion of its stomach content is, consequentially, impossible to identify through morphological examination (Asahida *et al.* 1997; Wilcox & Jeffries 1974). Furthermore, most studies on *C. crangon*'s diet to date have focused on a limited number of locations and relatively small spatial scales (e.g. Evans 1984; Oh *et al.* 2001; Pihl & Rosenberg 1984) while large scale studies are required to assess geographical variation in the shrimp's diet and to understand the relative importance of different prey items.

The degree to which food items are actively selected or passively ingested by consumers is an essential consideration in assessing the relative importance of different prey categories and understanding the trophic niche of consumers. Traditionally, indices are used to infer the predator's preference for prey based on the relative abundance of prey in the predator's diet and the prey's relative abundance in

the environment (e.g. Peterson & Ausubel 1984). Examples of commonly used indices are the Ivlev's Electivity Index (Ivlev 1961) and the Jacob's Index Of Selectivity (Jacobs 1974), which also corrects for item depletion (Jacobs 1974). Although some attempts have been made to link diet metabarcoding data with food availability in managed forests (Kowalczyk *et al.* 2011) and artificial mesocosms (Ray *et al.* 2016), no examples exist for wild marine animal trophic studies contrasted with whole-community environmental DNA data.

Here, we report on a large-scale analysis of the trophic ecology of *C. crangon*, which reveals its ecological role in estuarine systems and provides a key for the reconstruction of ecological networks of European coastal marine communities. By using nearly universal primers for mitochondrial cytochrome c oxidase I, we used metabarcoding to describe the diet of the shrimp, alongside the soft bottom communities on which they feed, over a European scale. We were expecting a wide variety of prey items, reflecting variation in environmental conditions and prey availability across European coasts. More specifically, we tested whether metabarcoding can (a) provide a detailed overview of *C. crangon's* diet, including prey selectivity, using DNA extracted from stomach and environmental samples; (b) identify geographical patterns in its trophic ecology, at both local and regional scales and (c) assess consistent and general trophic patterns in order to better define the ecological role of this widespread species.

Methods

Sample collection and processing

Brown shrimp and sediment samples were collected from 24 sites distributed over 6 estuaries in the Netherlands, Portugal and the United Kingdom (Fig. 1). Adult shrimp (>20 mm total length, TL; tip of the rostrum to tip of the telson) were captured in the intertidal zone (0-1m depth) by push-net at low tide (± 3 h). Shrimp (30-50 per site) were placed on ice and transported to the lab to be stored at -20 °C. Sediment was collected for the extraction of environmental DNA to characterise the biological community present at each site. Sediment was sampled from the upper 2 cm surface layer, which represent the most recent DNA deposits and the habitat where the shrimp live and feed (Pinn & Ansell 1993; Turner *et al.* 2015), with a PVC corer (3.2 mm \varnothing). Per site, three sediment subsamples were collected at several meters distance from each other and combined to reduce the influence of local heterogeneity (Taberlet *et al.* 2012b). The sediment was stored in 96 % ethanol, transported on ice and kept at -20 °C. At each site, temperature, salinity (Fisher Scientific Traceable Salinity Meter), pH (Hanna HI 98129), dissolved oxygen (OxyGuard Handy Mk I) and turbidity (Eutech TN-100) were measured in triplicates. Extra sediment was collected, in triplicates, from each site for granulometric analyses (Horiba LA-950 Particle size analyser) and Total Organic Matter (TOM) determination by means of ashing (550 °C, 6 h). One site (Mersey 3) was not included for analysis because high-throughput sequencing of its stomach samples did not result in sufficient read depth (< 1000 reads; see results).

DNA extraction

Overall, 1025 shrimp (20-50 mm TL) were caught and 494 full stomachs (visual determination) were dissected using flame-sterilised tools to avoid cross contamination. Stomachs were pooled in batches of 8 (from shrimp collected at the same site) prior to DNA extraction (Ray *et al.* 2016). Though the pooling of samples increases the number of stomachs analysed per sequencing run, pooling might

reduce the detected MOTU richness since large, recently ingested diet items in single stomachs may obscure the stomach contents of other individuals. Nevertheless, this potentially negative effect of pooling was deemed to be negligible, due to the pool replication conducted within location, and the population-emphasis of the study. Three replicate pools were extracted per site. However, due to a high percentage of empty stomachs in natural populations (20-60 %; Feller 2006; Oh *et al.* 2001; Pihl & Rosenberg 1984), some sites contained only 2 replicates and some replicates contained less than 8 full stomachs (see Supplementary table S1): the latter were still included in the analyses as variation in number of stomachs pooled did not affect the patterns observed (see results). In total, 66 pooled samples were extracted, divided over 24 sites. In addition to the full stomach samples, 3 pooled samples of 8 visually empty stomachs were included for comparative purposes.

DNA was extracted from 0.25 g of homogenized pooled stomach contents (N = 66) using the PowerSoil® DNA Isolation Kit (Mo-Bio laboratories), whereas DNA from sediment (10 g; N = 24) was extracted using the PowerMax® DNA Soil Kit (Mo-Bio laboratories). A Qubit fluorometer (Thermo Fisher Scientific) was used to assess DNA concentrations of purified extracts. DNA extraction and pre-PCR preparations were performed in separate labs from post-PCR procedures to reduce contamination.

DNA amplification and high-throughput sequencing

Amplification of DNA, for both stomach and sediment samples, was achieved using a single set of versatile, highly degenerated PCR primers targeting the 313-bp Leray fragment (Leray *et al.* 2013) of the mitochondrial cytochrome c. oxidase subunit I (COI) region. The mICOLintF-XT primer (5'-GGWACWRGWTGRACWITITAYCCYCC-3') was used as forward primer. This modified version (Wangensteen *et al.* 2018) of the mICOLintF primer (Leray *et al.* 2013) included 2 extra degenerate bases (equimolar mixtures of 2 different bases at a given position) and 2 inosine nucleotides (that can

match any nucleotide) to enhance its eukaryotic universality. The reverse primer was jgHCO2198 (5'-TAIACYTCIGGRTGICCRAARAAYCA-3'; Geller *et al.* 2013). The Leray fragment has already been successfully applied for both the characterisation of marine communities and marine fish gut contents (Leray & Knowlton 2015; Leray *et al.* 2015; Leray *et al.* 2013). Eight-base oligo-tags (Coissac *et al.* 2012) attached to the metabarcoding primers were added to the amplicons during a single PCR step, in order to label different samples in a multiplexed library. Also, a variable number (2, 3 or 4) of fully degenerate positions (Ns) was added at the beginning of each primer, in order to increase variability of the amplicon sequences (Guardiola *et al.* 2015). The PCR mix recipe included 10 µl AmpliTaq gold 360Master mix (Applied Biosystems), 3.2 µg Bovine Serum Albumin (Thermo Scientific), 1 µl of each of the 5 µM forward and reverse tagged-primers, 5.84 µl H₂O and 2 µl extracted DNA template (~ 5 ng µl⁻¹). The PCR profile included an initial denaturing step of 95 °C for 10 min, 35 cycles of 94 °C for 1 min, 45 °C for 1 min and 72 °C for 1 min and a final extension step of 72 °C for 5 minutes. After quality check of all amplicons by electrophoresis, the tagged PCR products (including 2 PCR negative controls) were pooled at equimolar concentration into 2 multiplexed sample pools (sediment and stomach) and purified using MinElute columns (Qiagen). Two Illumina libraries were subsequently built from these pools, using the NextFlex PCR-free library preparation kit (BIOO Scientific). Libraries were quantified using the NEBNext qPCR quantification kit (New England Biolabs) and pooled in a 1:4 sediment:stomach molar concentration ratio (similar to the sediment:stomach sample ratio) along with 0.7% PhiX (v3, Illumina) serving as a positive sequencing quality control. The libraries with a final molarity of 8 pM were sequenced on an Illumina MiSeq platform using v2 chemistry (2x250 bp paired-ends).

Preliminary analyses of the sequencing data revealed a substantial number of reads belonging to one Molecular Taxonomic Unit (MOTU) in the fungal order Hypocreales (Ascomycota). For further identification, the ITS fragment was amplified from 5 samples with a high number (>90% read abundance) of reads of this MOTU, with the primer combination ITS1f (5'-CTTGGTCATTTAGAGGAAGTAA-3'; Gardes & Bruns 1993) and ITS4ASCO (5'-

CGTTACTRRGGCAATCCCTGTTG-3'; Nikolcheva & Bärlocher 2004), specific for Ascomycota. The PCR mix recipe was the same as the one used for the Leray fragment described above and the PCR profile included an initial denaturing step of 95 °C for 5 min, 32 cycles of 95 °C for 30 sec, 55 °C for 30 sec and 72 °C for 1 min and a final extension step of 72 °C for 10 minutes (Manter & Vivanco 2007). After electrophoresis check, the amplicons of these 5 samples were cleaned and Sanger sequenced by Source Bioscience Sequencing UK.

Bioinformatic and data analyses

Bioinformatic analyses were performed using the OBITools metabarcoding software suite (Boyer *et al.* 2016). Read quality assessment was performed with FastQC and paired-end read alignment using illumina paired-end, retaining reads with an alignment quality score > 40. Demultiplexing and primer removal was achieved using ngsfilter with the default options. Obigrep was applied to select all aligned reads with a length between 303-323 bp and free of ambiguous bases. Obiuniq was used to dereplicate the reads and the uchime-denovo algorithm (Edgar *et al.* 2011) (implemented in VSEARCH; Rognes *et al.* 2016) was used to remove chimeras. Amplicon clustering was performed using the SWARM algorithm (Mahé *et al.* 2014, 2015) with a d value of 13, which offers a conservative solution to the high variability of the COI gene (Wangenstein & Turon 2017). After removal of singletons, taxonomic assignment of the representative sequences for each MOTU was performed using the ecotag algorithm (Boyer *et al.* 2016), using a local reference database (Wangenstein *et al.* 2018) containing filtered COI sequences retrieved from the BOLD database (Ratnasingham & Hebert 2007) and the EMBL repository (Kulikova *et al.* 2004). This algorithm uses a phylogenetic approach to assign sequences to the most reliable monophyletic unit, based on the density of the reference database. The data was refined by clustering MOTUs assigned to the same species, abundance renormalization (to remove false positives due to tag-switching; Wangenstein & Turon 2017) and by removing bacterial reads and contaminations of human or terrestrial origin. MOTUs with a maximum of four or less reads per sample were removed on a sample by sample basis to avoid false positives and low

frequency noise (De Barba *et al.* 2014; Wangensteen *et al.* 2018). All MOTUs for which the abundance in the PCR negative controls was higher than 10 % of the total reads of that MOTU were removed (Wangensteen & Turon 2017). Samples with a low read depth (< 1000) following removal of predator, parasite and contaminant reads were removed prior to analysis. All statistical analyses were performed in R v3.1.3 (<https://www.R-project.org/>) with the vegan (v2.3-5) and BiodiversityR (v2.5-3) packages (Kindt & Coe 2005; Oksanen *et al.* 2016). Only MOTUs showing mean relative abundance \geq 0.5% in the full stomach samples were considered (Albaina *et al.* 2016) for non-metric multidimensional scaling (nMDS), canonical correspondence analysis (CCA) and PERMANOVA analyses. Correlation between sediment and stomach community composition was tested with a Mantel test (Bray-Curtis dissimilarities; Pearson's product-moment correlation; 999 permutations). The influence of environmental variables (mean temperature, salinity, pH, oxygen saturation, turbidity, median sediment grain size and TOM) on the full stomach contents were tested by means of CCA and PERMANOVA. PERMANOVAs were calculated using the function Adonis (vegan) with Bray-Curtis dissimilarities and 1000 permutations. Prior to CCA and PERMANOVA, model selection was performed using the function ordistep (vegan). Prey MOTU richness for each estuary was represented as MOTU accumulation curves after rarefaction for the number of reads (1000 reads, 500 permutations) and the number of samples (9-15 samples, 1000 permutations). The Jacobs' Selectivity Index was calculated based on the relative read abundances of the MOTUs extracted from sediment and stomach samples in accordance to Jacobs (1974). Trophic significance of individual MOTU was determined based on the relative read abundance, fraction of samples with MOTU presence and Jacobs' Selectivity Index as follows: Trophic significance = (relative abundance) * (fraction of samples) * (Jacobs' Selectivity Index + 1). Trophic significance was represented in categorical terms based on the relative trophic significance of each MOTU (high: > 10 %, medium 1 % - 9 %, low < 1 %) instead of exact values since the relative abundances of individual taxa should be considered with caution (Deagle *et al.* 2005).

Results

Collection statistics

A total of 1025 *C. crangon* were caught with a 1:8 male:female sex ratio (based on 767 shrimp which could be sexed morphologically). About 7.5 % of the females were ovigerous. Mean (\pm SD) wet weight was 0.40 ± 0.26 gram; mean (\pm SD) TL was 35.1 ± 7.6 mm and mean (\pm SD) carapace length (CL) was 7.4 ± 1.6 mm ($CL = 0.214 * TL$; $r^2 = 0.81$, $N = 1025$). TL varied significantly between sites (Supplementary table S2; One-way ANOVA: $Df = 23$, $F = 47.95$, $P < 0.001$). Overall, the proportion of *C. crangon* with a full stomach was 57.9%. Mean proportion of full stomachs per site ($58.9 \pm 19.3\%$) was not correlated with the time of sampling (Pearson's correlation: $R^2 = 0.07$, $P = 0.754$, $N = 24$).

High-throughput DNA sequencing

A total number of 8,895,448 reads were obtained from an Illumina MiSeq run of pooled amplicon libraries built from 24 sediment samples, 69 pooled *C. crangon* full stomach samples (from now on referred to as stomach samples), 3 pooled *C. crangon* empty stomach samples (comprising of stomach tissue and clear liquid) and 2 PCR negative controls. Variation in the number of pooled stomachs did not affect the patterns of diet composition (PERMANOVA: pseudo- $F_{1,50} = 1.0$, $P = 0.453$) or MOTU richness (rarefied to 1000 reads) per sample (generalized linear model with quasipoisson distribution: $t = 1.08$, $P = 0.650$). In total, 5,704,471 reads remained after sample demultiplexing, quality and sequence-length filtering, and removal of bacterial reads, contaminations and false-positives due to tag-switching (sediment samples: 742,286; stomach samples: 4,828,136; empty stomach samples: 134,049). After taxonomic assignment, a total of 39 MOTUs (16 Metazoa, 7 Rhodophyta, 5 Stramenopiles, 11 unassigned) were removed because their abundance in the PCR negative control was $> 10\%$ of the total reads of those MOTUs. Mean (\pm SD) proportion of *C. crangon* reads was $28 \pm 29\%$ (range: 0.2-97.6 %) in the stomach samples and $47 \pm 46\%$ (range: 10.6-99.1 %) in the empty

271 stomach samples. Mean proportion of *C. crangon* reads was 1 ± 4 % in the sediment samples (range:
 272 0.0-21.0 %). Remaining number of reads per sample ranged 179-203,808 in full stomach, 7-332 in
 273 empty stomach and 5,114-71,770 in sediment samples. A high number of reads (4,828,136 reads)
 274 belonging to a fungus of the species *Purpureocillium lilacinum* (Ascomicota: Hypocreales) was
 275 detected in almost all (95 %) stomach samples and identified using both COI (100 % identity) and ITS
 276 markers (100 % identity; Supplementary table S3). Mean (\pm SD) proportion of *P. lilacinum* reads was
 277 36 ± 37 % (range: 0.0-97.4 %) in full stomach, 53 ± 47 % (range: 0.1-89.4 %) in empty stomach and 0.1
 278 ± 0.2 % (range: 0.0-0.8 %) in sediment samples. No *P. lilacinum* were detected in the PCR negative
 279 controls. All *C. crangon* and *P. lilacinum* reads were removed from the database prior to further
 280 analyses on diet, resulting in a total of 2,687,877 reads divided over 66 pooled stomach samples (Fig.
 281 2) and 24 sediment samples (Fig. 3). A total of 14 pooled stomach samples were removed, prior to
 282 further analyses, because they contained less than 1000 diet-related reads (Supplementary table S4).
 283 One sediment sample (Mersey 3) was, consequently, also removed since no stomach samples were
 284 included for that site. The final dataset consisted of a total of 8,321 MOTUs, of which 6,299 MOTUs
 285 belonging to 40 phyla were detected in the sediment samples, 2,342 (35 phyla) in the stomach
 286 samples, and 14 (7 phyla) in the empty stomach samples. A total of 502 MOTUs were detected both
 287 in the sediment and stomach samples and only 2 (an unassigned Rhodophyta and an unassigned
 288 Eukaryota) were detected exclusively in the empty stomach samples. Of the total number of MOTUs
 289 detected, 370 could be assigned to the species level of which 291 were detected in the stomach
 290 samples. Twenty taxa showed a relative abundance greater than 5 % in any given sample while 85 taxa
 291 showed an abundance greater than 1%. The final number of diet-related reads per stomach sample
 292 varied randomly, without systematic trends across estuaries (One-way ANOVA: Estuary: $F_{5,29} = 2.017$,
 293 $P = 0.106$; Sites nested in estuary: $F_{17,29} = 0.811$, $P = 0.669$). Rarefaction curves (Supplementary Fig.
 294 S1) showed that a plateau in the number of MOTUs was achieved in almost all cases, indicating an
 295 overall sequencing depth adequate to capture the number of MOTUs present. Empty stomach

samples contained a very low number of MOTUs and reads, and were, therefore, not taken into account for any further analyses (Fig. 4A).

Description of *Crangon crangon* diet

Analysis of *C. crangon* stomach contents showed large variation in relative MOTU abundances between samples (Fig. 2). Notable patterns are: the lack of a dominant MOTU detected in stomachs from the Aveiro estuary; a relatively high (10-25 %) contribution of the decapod crabs *Carcinus maenas* and *Pisidia longicornis* in the Minho estuary; the detection of the introduced barnacle *Austrominius modestus* in the Scheldt and Mersey estuaries; high amounts (~25 %) of the polychaete *Pista cristata* in the Eastern Scheldt; the substantial proportion (~10 %) of the mysid *Neomysis integer* reads in the Mersey estuary, and the dipteran *Chironomus salinarius* also, in the most inland sampling site of this estuary (> 75 %); the large contribution (~ 50 %) of the amphipod *Corophium volutator* in the Kent estuary. In general, the shore crab *C. maenas* and the amphipod *C. volutator* were the trophically preponderant prey items for *C. crangon* (Table 1). Other important MOTUs included annelids (*Hediste diversicolor* and *P. cristata*), other amphipods (*Bathyporeia sarsi*), other decapods (*P. longicornis*), chironomids (unassigned), mysids (*N. integer*) and barnacles (*A. modestus*). Fish reads were detected in all estuaries with a total of 22 species present in 27 stomach samples. Five fish species were relatively abundant (≥ 5 %; Table 1) but were generally only present in a small number of stomach samples. One other noteworthy observation is the presence of low abundances of known parasitic taxa in several stomach samples (*Hematodinium* sp., 3 stomach samples; Apicomplexa, 6 stomach samples).

Selectivity in *Crangon crangon* diet

MOTU diversity within phyla was generally higher in the sediment than in the stomach samples, except for Arthropoda, Annelida, Mollusca and Chordata (Fig. 4A). The proportion of MOTUs that could not be assigned at the phylum level was higher in the sediment (73 %) than in the stomach samples (58 %) and many abundant taxa in the sediment could not be identified at lower taxonomic ranks (Fig. 3). Data combined per sample type (sediment/stomach) and MOTUs pooled at the phylum level showed that sediment samples contained high relative read abundances of Bacillariophyta (20 ± 3 %), Discosea (10 ± 2 %), Dinoflagellata (6 ± 2 %) and Arthropoda (5 ± 1 %) while *C. crangon* stomach samples contained a high mean (\pm SE) relative read abundance (%) for Arthropoda (53 ± 5 %), Annelida (12 ± 3 %) and Chordata (5 ± 2 %; Fig. 4B). Mantel test results showed a significant correlation between the community structure detected in the stomach and sediment samples ($r: 0.43$, $P < 0.01$), indicating an association between the shrimp's diet and its prey abundance in the environment. Analyses of DNA extracted from both sediment and *C. crangon* pooled stomach samples showed, based on all MOTUs detected, significant differences between sample types and estuaries (Fig. 5 and 6A; PERMANOVA: sample type: pseudo- $F_{1,68} = 7.8$, $P < 0.001$; estuary: pseudo- $F_{5,68} = 2.5$, $P < 0.001$). Visual inspection of the relative abundances of the most important MOTUs also showed a high discrepancy between the abundances in the stomach (Fig. 2) and sediment samples (Fig. 3). MOTUs abundant in the stomach samples (≥ 1 % abundance) showed, furthermore, a low read abundance in the sediment samples in all estuaries (Fig. 6A). These differences in relative abundances, resulted in many MOTUs having a maximum Jacobs' selectivity index value of 1 which indicates that prey items were highly selected (Table 1). Phylum composition differed significantly between sediment and full stomach samples (PERMANOVA: pseudo- $F_{1,44} = 34.1$, $P < 0.001$). Apart from Cnidaria and Rhodophyta, all phyla with ≥ 1 % abundance in either sediment or stomach samples showed significant differences (based on paired Wilcoxon signed-rank tests) in relative read abundances between the sediment and stomach samples (Supplementary table S5). Visualisation of the importance of the phyla detected in the stomach samples based on the mean relative abundance (%), presence (%) and Jacobs' selectivity index (D) is shown in Fig. 4C.

Variation between estuaries

Multivariate analysis on the stomach contents (MOTUS ≥ 0.5 % abundance) showed significant differences between estuaries (Fig. 6B; PERMANOVA: pseudo- $F_{5,29} = 2.7$, $P < 0.001$) and sites nested within estuaries (PERMANOVA: pseudo- $F_{17,29} = 0.6$, $P < 0.001$). Bonferroni-corrected pairwise comparisons showed similarity in consumed community structure among the Eastern Scheldt, Western Scheldt and Mersey estuaries. Minho differed significantly from the Mersey and Eastern Scheldt estuaries. Aveiro or Kent significantly differed from all other estuaries (See Supplementary table S6 for details). Step-wise model selection (both forward and reverse) and CCA (Fig. 6C) showed significant influences of salinity ($P < 0.01$), median grain size ($P < 0.01$) and TOM ($P < 0.05$; see table S7 for means per estuary) on MOTU composition in *C. crangon* stomach samples (≥ 0.5 % abundant MOTUs). The environmental variables (constrained CCA axes) explained 34 % of the variance in the dataset. Temperature, turbidity and oxygen saturation did not have a significant influence on the model and pH was strongly correlated with salinity ($r^2 = 0.75$, $P < 0.001$, $N = 23$). These factors were, therefore, not included in the final model. MOTU richness (rarefied to 1000 reads) in *C. crangon* stomach contents also showed differences between estuaries, with the Aveiro and Eastern Scheldt estuaries showing a higher number of MOTUs than the others (Fig. 7). The slopes of the MOTU accumulation curves, however, did not approach an asymptote, offering a glimpse of the vast amount of marine biodiversity yet to be uncovered.

Discussion

Evaluation of *C. crangon* diet

This study provides a detailed overview of the brown shrimp's trophic ecology, focusing on dietary variations at multiple geographical scales. Adult brown shrimp were caught in a variety of sandy estuarine intertidal habitats. Mainly females were captured, probably due to the spatial sex-specific segregation of *C. crangon* during the summer-autumn period (Bamber & Henderson 1994; Henderson & Holmes 1987). The results confirm *C. crangon* as a generalist consumer feeding on a broad variety of food items but preferring larger mobile epifaunal prey items such as crustaceans, annelids and fish. The present investigation uncovered a great diet contribution of decapods and teleosts, while these were usually not considered to be important contributors to the shrimp diet in previous studies (e.g. Ansell *et al.* 1999; Oh *et al.* 2001; Plagmann 1939; Raffaelli *et al.* 1989). Although comparisons of prey contribution should be made cautiously due to diversity of quantification methods used, the observed trend could be partly explained by scavenging behaviour on large organisms, previously not recorded in crangonid shrimps. Crangonid shrimps generally macerate and eat the soft body parts of larger preys (Asahida *et al.* 1997; Gibson *et al.* 1995; Seikai *et al.* 1993; Wilcox & Jeffries 1974). Smaller food items, on the other hand, are often ingested as a whole, including their hard body parts (Tiews 1970) and are thus more easily identified by morphological methods. This discrepancy in detectability might possibly have played a role in studies that have detected low amount of fish and decapods but considerable amounts of unidentified soft tissue (e.g. Oh *et al.* 2001; Raffaelli *et al.* 1989). Metabarcoding methods can detect and taxonomically identify such soft tissues, thus highlighting the enhanced suitability of molecular approaches to present a more realistic picture of trophic ecology in marine invertebrates.

The diet of *C. crangon* showed a high MOTU richness, including previously described food items (Table 1). The number of COI MOTUs (2,342) detected in the shrimp's stomachs may be an overestimation

of the total number of real species (e.g. due to detection of pseudogenes; Tang *et al.* 2012; Vamos *et al.* 2017) and includes protists and microalgae (Wangensteen *et al.* 2018), which are unlikely to be prey items of *C. crangon*. Nonetheless, even just the 306 ascertained or 20 most abundant (> 0.5 % relative abundance) species in the shrimp's diet was remarkably higher than the number found in previous studies based on morphological identification (see Table 1). Furthermore, twenty taxa showed a high abundance (>5 %) in any given sample, probably representing important prey items at some locations or times.

Two species were predominant in our study: the shore crab *Carcinus maenas* across the overall geographic distribution, and the amphipod *Corophium volutator* in UK localities (characterised by muddy sediments and high organic matter content). Both species are well-known prey of *C. crangon* (Evans 1984; Moksnes *et al.* 1998; Pihl & Rosenberg 1984) and can occur at high densities in soft-bottom estuarine habitats (Meadows & Reid 1966; Moksnes 2002). Consumption of *C. maenas* could be the result of scavenging, although juvenile crabs could be captured, while *C. volutator* are likely to be predated, as these amphipods are small (up to 11 mm TL). Overall, the local distribution of the detected food items followed environmental gradients reflecting their ecology. Euryhaline deposit feeders such as *C. volutator* and *H. diversicolor*, *N. integer* and *Chironomus salinarius* (larval stage) were mainly associated with muddy, brackish sites with high organic matter content, commonly inhabited by these species (Anderson 1972; Drake & Arias 1995; Mauchline 1971; Meadows 1964; Mees *et al.* 1993; Ólafsson & Persson 1986). Stomach samples taken from sites with larger grain size contained species adapted to coarser sands, such as *P. longicornis* and *Talitrus saltator* (Fanini *et al.* 2007; Pallas *et al.* 2006). Detection of fish DNA could reflect a combination of direct predation on juveniles of species which use the estuaries as nurseries (e.g., *Platichthys flesus* and *Dicentrarchus labrax*) and scavenging on dead bodies of species which do not regularly use estuaries as a nursery (e.g., *Scomber scombrus* and *Labrus bergylta*; Elliott & Dewailly 1995). The high presence of the

invasive barnacle *A. modestus* DNA at several locations was likely due to the capture of cypris or nauplii larvae (Ansell *et al.* 1999; Boddeke *et al.* 1986).

This is also the first study showing a high occurrence of *P. lilacinum* (Ascomycota: Hypocreales) in the digestive system of *C. crangon*. *Purpureocillium lilacinum* is a well-studied fungus, being abundant in terrestrial soils (Cham Thi Mai *et al.* 2016) and detected in the marine environment (Redou *et al.* 2015; Yue *et al.* 2015). It is a known pathogen of nematodes and therefore of commercial importance as a biological control agent to manage pests of several crops (Castillo Lopez *et al.* 2014; Singh *et al.* 2013). This fungus is even considered to be of medical importance since it can infect humans and other vertebrates with compromised immune systems (Luangsa-Ard *et al.* 2011). As *P. lilacinum* has been successfully cultured (for the production of chitosanase) using farmed marine shrimp by-products as substrate (*Penaeus* sp.; Nidheesh *et al.* 2015) and is closely related to known parasites of crabs (Smith *et al.* 2013), it might be postulated that it has a symbiotic relationship with *C. crangon*, although more research is required to test this hypothesis. Its occurrence and high relative abundance (although possibly overestimated since its DNA was extracted from a living community, as opposed to digested food) in *C. crangon* stomach samples over a large geographical area are clear indicators that this species might be important for the brown shrimp's ecology and/or physiology. Alongside *P. lilacinum*, several other known parasitic taxa have been detected in the shrimp's stomachs, including *Hematodinium* sp. and Apicomplexa (Molnar *et al.* 2012; Rueckert *et al.* 2011; Stentiford & Shields 2005).

The application of metabarcoding in crustacean trophic studies

Metabarcoding using universal primers is generally considered as a simple, rapid and relatively inexpensive method to define in detail the feeding ecology of organisms (Berry *et al.* 2015; Kartzinel

& Pringle 2015; Pinol *et al.* 2014). The fraction of the brown shrimp DNA detected in its own gut was low allowing for the detection of prey items without using predator-specific blocking primers (average: 28%; compared to e.g. Olmos-Perez *et al.* 2017; Pinol *et al.* 2014). Metabarcoding has several clear advantages over traditional trophic methods including the better detection of soft-bodied, small and cryptic taxa, higher speed of analysis (Berry *et al.* 2015; Casper *et al.* 2007; Chariton *et al.* 2015; Symondson 2002), and traceability of identifications, which do not rely on the availability of morphological taxonomic expertise. Furthermore, the application of metabarcoding even allows for the detection of prey items in empty guts (Harms-Tuohy *et al.* 2016), albeit the DNA extracted from visually empty *C. crangon* stomachs was too low in prey read number and MOTU richness for robust comparisons.

Both traditional morphological examination and DNA-metabarcoding of food items suffer from limitations in providing quantitative descriptions of the diet of consumers (Casper *et al.* 2007). For metabarcoding, errors can occur due to technical artefacts specific to DNA amplification and sequencing (Barnes & Turner 2016; Pompanon *et al.* 2012), and biological limitations such as species-specific digestion and DNA degradation rates (Deagle *et al.* 2010; Murray *et al.* 2011; Pinol *et al.* 2014; Sakaguchi *et al.* 2017). Furthermore, some of the DNA detected might come from secondary predation (taxa present in the stomach of preyed organisms; Berry *et al.* 2015; Kartzinel & Pringle 2015). Cannibalism also imposes a specific problem in trophic molecular studies since it cannot be identified by means of metabarcoding (Berry *et al.* 2015; Ray *et al.* 2016). Large brown shrimps are known to be cannibalistic (Evans 1984; Pihl & Rosenberg 1984) but the removal of *C. crangon* sequence reads from our data set makes it impossible to gauge insights into the extent of cannibalism in this species. Due to the restrictions in the quantification of consumed prey volume, many trophic studies only use presence/absence data (e.g. Deagle *et al.* 2010; Harms-Tuohy *et al.* 2016; Pinol *et al.* 2015). This might, however, result in an overestimation of small taxa that are abundant in the sediment, but with low

464 trophic relevance, as they could, in the case of *C. crangon*, be passively acquired when shrimp ingest
465 sediment to crush food in their stomach (Ansell *et al.* 1999; Deagle *et al.* 2018; Tiews 1970). Multiple
466 stomachs were pooled prior to analysis and data was subjected to rigorous filtering to allow for a semi-
467 quantitative estimation of proportions of prey DNA (Deagle *et al.* 2005; Lejzerowicz *et al.* 2015;
468 Pompanon *et al.* 2012; Thomas *et al.* 2016). Relative abundances of individual taxa should, however,
469 be considered with caution and viewed more in categorical terms (low or high trophic significance)
470 than exact proportions (Deagle *et al.* 2005). This study provides a significant addition to a growing
471 body of studies in showing the applicability of semi-quantitative estimations in molecular trophic
472 ecology (e.g. Albaina *et al.* 2016; Deagle *et al.* 2018; Ray *et al.* 2016; Sakaguchi *et al.* 2017; Soininen *et*
473 *al.* 2013).

474
475 Finally, the results presented draw a close link between prey distribution in estuarine habitats and
476 ingested prey item abundance. The use of eDNA from sediments to assess community composition
477 and to relate this to the shrimp's diet is a novel contribution to the fields of molecular trophic analysis
478 and eDNA, which goes beyond the taxon studied. It should be noted, however, that a correct
479 assessment of the predator's trophic niche by means of prey selectivity determination relies on a
480 correct assessment of prey abundance, both in the stomach and in the environment. Issues with
481 incorrect abundance estimations, for example due to species-specific detection rates (e.g. due to
482 different rates of DNA sequestering in the environment; Barnes & Turner 2016), are not specific to
483 molecular studies (Strauss 1979) and the constant work on improving the reliability of relative
484 abundance estimations from eDNA (Deagle *et al.* 2018; Thomas *et al.* 2016; Ushio *et al.* 2018) should
485 substantially enhance the applicability of selectivity indices in molecular research.

486
487 **Geographic variation in *C. crangon* trophic ecology**

This study also assesses for the first time a large geographical variation in the brown shrimp's trophic ecology at multiple spatial scales. Previous studies have shown local variability in *C. crangon* diet (Evans 1984; Oh *et al.* 2001; Pihl & Rosenberg 1984) but no studies have been performed across multiple European estuaries. The results indicate that the consumed prey community can vary at local (within estuary, as discussed above) and regional (between estuaries) scales. The seasonal and tidal migratory behaviour of *C. crangon* (Al-Adhub & Naylor 1975; Henderson & Holmes 1987) may complicate localized diet assessments since their stomach contents might also contain food consumed at distant locations. Yet, this effect is considered to be minimal since the brown shrimp's relatively fast gut passage time ensures that their stomach contents mainly contain recently consumed items (4-20h; Feller 2006; Pihl & Rosenberg 1984; van der Veer & Bergman 1987). Large scale assessment of *C. crangon*'s trophic ecology showed high similarity between the Eastern Scheldt, Western Scheldt and Mersey estuaries and distinct diets in the Aveiro, Minho and Kent estuaries. The Aveiro estuary forms a large, saline lagoon with a wide variety of different habitats incorporating euhaline/polyhaline areas with relatively high species richness (Rodrigues *et al.* 2011). On the other hand, the Minho estuary is characterized by high water discharge, salinity variations (Costa-Dias *et al.* 2010) and significantly larger sediment grain size, factors which determine significantly divergent biodiversity features. The Kent estuary has a low species diversity caused by its fine sediments and low salinity (Anderson 1972). The Mersey estuary also showed a relatively low species richness detected in the stomach contents of *C. crangon*, probably related to its history of anthropogenic stress (Jones 2000). Overall, trophic variation in *C. crangon* depends on patterns in the local abundance and distribution of its prey (in line with: Oh *et al.* 2001; Pihl 1985; Pihl & Rosenberg 1984). In order to evaluate this variation more exhaustively, knowledge on the ecology and seasonality of the local macrozoobenthic community is required.

***Crangon crangon*'s ecological role**

Based on the results of this study, *C. crangon* can best be described as a highly opportunistic carnivore and scavenger. Despite its broad dietary range (Fig. 8), it shows, nevertheless, a prominent level of selectivity to larger motile mobile epifaunal prey items. This high level of flexibility in its trophic ecology might contribute to its very wide distribution on European coasts (Campos *et al.* 2009). In order to feed on diverse prey taxa, adult *C. crangon* are capable of employing a variety of methods (Fig. 8) including ambush predation (Gibson *et al.* 1995; Pinn & Ansell 1993), gulping behaviour (Tiews 1970) and scavenging (Fig. 8; Ansell *et al.* 1999; Price 1962). Since meiofaunal and protist phyla were not selected as prey items based on Jacobs' selectivity index (present but not abundant; in line with: Evans 1983; Feller 2006), it is possible that these taxa were passively consumed during the ingestion of sand to aid digestion (Ansell *et al.* 1999; Tiews 1970) or through secondary predation. Several studies classify *C. crangon* as an omnivore (Ansell *et al.* 1999; Raffaelli *et al.* 1989; Tiews 1970), but we cannot confirm this classification, because the primers used during this study have a very low affinity for chlorophytes resulting in many algal taxa not being detected (Wangensteen *et al.* 2018). Nevertheless, the algal phyla that can be detected with these primers (e.g. Rhodophyta, Phaeophyta and Bacillariophyta) had a low selectivity, indicating a negligible trophic importance for *C. crangon*. More research is required with plant-specific primers to assess the actual contribution of herbivory to the diet of the brown shrimp. Overall, the results of this study yield a level understanding of the trophic ecology of this species that would not have been possible through traditional morphological analysis, and which is key to providing essential insights into coastal community interactions.

Acknowledgements

We would like to express our thanks to C. Adiele, A. Bio, T. Ferraz, R. Fraser, A. Goossens, P. Oldfield and E. Siegenthaler for their help collecting shrimp under all weather conditions; C. Hide and M. Patel for their practical assistance in the lab and A. Mastin and E. Grist for their advice on data analysis. We are also grateful to J. Taylor for his advice regarding the amplification of fungal DNA and

538 I. Goodhead for his advice on Illumina sequencing. Funding for this project was provided by the
539 University of Salford, Mersey Gateway Environmental Trust, the Santander Advantage fund, and the
540 Natural Environment Research Council, grant NE/N005759/1.

541

References

- Al-Adhub AHY, Naylor E (1975) Emergence rhythms and tidal migrations in the brown shrimp *Crangon crangon* (L.). *Journal of the Marine Biological Association of the United Kingdom* **55**, 801-810.
- Albaina A, Aguirre M, Abad D, Santos M, Estonba A (2016) 18S rRNA V9 metabarcoding for diet characterization: a critical evaluation with two sympatric zooplanktivorous fish species. *Ecology and Evolution* **6**, 1809-1824.
- Anderson SS (1972) The ecology of Morecambe Bay. II. Intertidal invertebrates and factors affecting their distribution. *Journal of Applied Ecology* **9**, 161-178.
- Ansell AD, Comely CA, Robb L (1999) Distribution, movements and diet of macrocrustaceans on a Scottish sandy beach with particular reference to predation on juvenile fishes. *Marine Ecology Progress Series* **176**, 115-130.
- Asahida T, Yamashita Y, Kobayashi T (1997) Identification of consumed stone flounder, *Kareius bicoloratus* (Basilewsky), from the stomach contents of sand shrimp, *Crangon affinis* (De Haan) using mitochondrial DNA analysis. *Journal of Experimental Marine Biology and Ecology* **217**, 153-163.
- Aviat D, Diamantis C, Neudecker T, Berkenhagen J, Müller M (2011) The north sea brown shrimp fisheries, pp. 5-103. European Parliament, Policy Department B: Structural and Cohesion Policies, Brussels.
- Bamber RN, Henderson Pa (1994) Seasonality of caridean decapod and mysid distribution and movements within the Severn Estuary and Bristol Channel. *Biological Journal of the Linnean Society*, 83-91.
- Barnes MA, Turner CR (2016) The ecology of environmental DNA and implications for conservation genetics. *Conservation Genetics* **17**, 1-17.
- Berry O, Bulman C, Bunce M, *et al.* (2015) Comparison of morphological and DNA metabarcoding analyses of diets in exploited marine fishes. *Marine Ecology Progress Series* **540**, 167-181.
- Boddeke R, Driessen G, Doesburg W, Ramaekers G (1986) Food availability and predator presence in a coastal nursery area of the brown shrimp (*Crangon crangon*). *Ophelia* **26**, 77-90.
- Boyer F, Mercier C, Bonin A, *et al.* (2016) OBITOOLS: a unix-inspired software package for DNA metabarcoding. *Molecular Ecology Resources* **16**, 176-182.
- Campos J, Pedrosa C, Rodrigues J, *et al.* (2009) Population zoogeography of brown shrimp *Crangon crangon* along its distributional range based on morphometric characters. *Journal of the Marine Biological Association of the United Kingdom* **89**, 499-499.
- Campos J, van der Veer HW (2008) Autecology of *Crangon crangon* (L.) with an emphasis on latitudinal trends. *Oceanography and Marine Biology: An Annual Review* **46**, 65-104.
- Casper RM, Jarman SN, Deagle BE, Gales NJ, Hindell MA (2007) Detecting prey from DNA in predator scats: A comparison with morphological analysis, using *Arctocephalus* seals fed a known diet. *Journal of Experimental Marine Biology and Ecology* **347**, 144-154.
- Castillo Lopez D, Zhu-Salzman K, Ek-Ramos MJ, Sword GA (2014) The entomopathogenic fungal endophytes *Purpureocillium lilacinum* (formerly *Paecilomyces lilacinus*) and *Beauveria bassiana* negatively affect cotton aphid reproduction under both greenhouse and field conditions. *PLoS One* **9**, e103891.
- Cham Thi Mai L, Nhi Thi Thuy L, Duong Thi Thuy N, Hoang Nguyen Duc P, Xo Hoa D (2016) The genus *Purpureocillium* from different ecology in the Southeast Vietnam. *International Journal of Agricultural Technology* **12**, 2255-2274.
- Chariton AA, Stephenson S, Morgan MJ, *et al.* (2015) Metabarcoding of benthic eukaryote communities predicts the ecological condition of estuaries. *Environmental Pollution* **203**, 165-174.
- Coissac E, Riaz T, Puillandre N (2012) Bioinformatic challenges for DNA metabarcoding of plants and animals. *Molecular Ecology* **21**, 1834-1847.

- Costa-Dias S, Freitas V, Sousa R, Antunes C (2010) Factors influencing epibenthic assemblages in the Minho Estuary (NW Iberian Peninsula). *Marine Pollution Bulletin* **61**, 240-246.
- De Barba M, Miquel C, Boyer F, *et al.* (2014) DNA metabarcoding multiplexing and validation of data accuracy for diet assessment: application to omnivorous diet. *Molecular Ecology Resources* **14**, 306-323.
- Deagle BE, Chiaradia A, McInnes J, Jarman SN (2010) Pyrosequencing faecal DNA to determine diet of little penguins: is what goes in what comes out? *Conservation Genetics* **11**, 2039-2048.
- Deagle BE, Thomas AC, McInnes JC, *et al.* (2018) Counting with DNA in metabarcoding studies: How should we convert sequence reads to dietary data? *Molecular Ecology* **00**, 1-16.
- Deagle BE, Tollit DJ, Jarman SN, *et al.* (2005) Molecular scatology as a tool to study diet: analysis of prey DNA in scats from captive Steller sea lions. *Molecular Ecology* **14**, 1831-1842.
- Drake P, Arias AM (1995) Distribution and production of *Chironomus salinarius* (Diptera: Chironomidae) in a shallow coastal lagoon in the Bay of Cádiz. *Hydrobiologia* **299**, 195-206.
- Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R (2011) UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* **27**.
- Elliott M, Dewailly F (1995) The structure and components of European estuarine fish assemblages. *Netherlands Journal of Aquatic Ecology* **29**, 397-417.
- Evans S (1983) Production, predation and food niche segregation in a marine shallow soft-bottom community. *Marine Ecology Progress Series* **10**, 147-157.
- Evans S (1984) Energy budgets and predation impact of dominant epibenthic carnivores on a shallow soft bottom community at the Swedish west coast. *Estuarine, Coastal and Shelf Science* **18**, 651-672.
- Fanini L, Marchetti GM, Scapini F, Defeo O (2007) Abundance and orientation responses of the sandhopper *Talitrus saltator* to beach nourishment and groynes building at San Rossore natural park, Tuscany, Italy. *Marine Biology* **152**, 1169-1179.
- Feller RJ (2006) Weak meiofaunal trophic linkages in *Crangon crangon* and *Carcinus maenus*. *Journal of Experimental Marine Biology and Ecology* **330**, 274-283.
- Gardes M, Bruns TD (1993) ITS primers with enhanced specificity for basidiomycetes - application to the identification of mycorrhizae and rusts. *Molecular Ecology* **2**, 113-118.
- Geller J, Meyer C, Parker M, Hawk H (2013) Redesign of PCR primers for mitochondrial cytochrome c oxidase subunit I for marine invertebrates and application in all-taxa biotic surveys. *Molecular Ecology Resources* **13**, 851-861.
- Gibson RN, Yin MC, Robb L (1995) The behavioural basis of predator-prey size relationships between shrimp (*Crangon crangon*) and juvenile plaice (*Pleuronectes platessa*). *Journal of the Marine Biological Association of the United Kingdom* **75**, 337-349.
- Guardiola M, Uriz MJ, Taberlet P, *et al.* (2015) Deep-sea, deep-sequencing: Metabarcoding extracellular DNA from sediments of marine canyons. *PLoS One* **10**, e0139633.
- Hanski I, Hansson L, Henttonen H (1991) Specialist predators, generalist predators, and the microtine rodent cycle. *The Journal of Animal Ecology* **60**, 353.
- Harms-Tuohy CA, Schizas NV, Appeldoorn RS (2016) Use of DNA metabarcoding for stomach content analysis in the invasive lionfish *Pterois volitans* in Puerto Rico. *Marine Ecology Progress Series* **558**, 181-191.
- Henderson PA, Holmes RHA (1987) On the population biology of the common shrimp *Crangon crangon* (L.) (Crustacea: Caridea) in the Severn Estuary and Bristol Channel. *Journal of the Marine Biological Association of the United Kingdom* **67**, 825-847.
- Holling CS (1965) The functional response of predators to prey density and its role in mimicry and population regulation. *Memoirs of the Entomological Society of Canada* **97**, 5-60.
- Ivlev VS (1961) *Experimental ecology of the feeding of fishes*. Yale University Press, New Haven, Connecticut, USA.
- Jacobs J (1974) Quantitative measurement of food selection. *Oecologia* **14**, 413-417.

- Jones PD (2000) The Mersey estuary – back from the dead? Solving a 150-year old problem. *Water and Environment Journal* **14**, 124-130.
- Kartzinel TR, Pringle RM (2015) Molecular detection of invertebrate prey in vertebrate diets: trophic ecology of Caribbean island lizards. *Molecular Ecology Resources* **15**, 903-914.
- Kindt R, Coe R (2005) *Tree diversity analysis: a manual and software for common statistical methods for ecological and biodiversity studies* World Agroforestry Centre, Nairobi (Kenya).
- Kowalczyk R, Taberlet P, Coissac E, *et al.* (2011) Influence of management practices on large herbivore diet—Case of European bison in Białowieża Primeval Forest (Poland). *Forest Ecology and Management* **261**, 821-828.
- Kulikova T, Aldebert P, Althorpe N, *et al.* (2004) The EMBL nucleotide sequence database. *Nucleic Acids Research* **32**, D27-D30.
- Lasley-Rasher RS, Brady DC, Smith BE, Jumars PA (2015) It takes guts to locate elusive crustacean prey. *Marine Ecology Progress Series* **538**, 1-12.
- Lejzerowicz F, Esling P, Pillet L, *et al.* (2015) High-throughput sequencing and morphology perform equally well for benthic monitoring of marine ecosystems. *Scientific Reports* **5**, 13932.
- Leray M, Knowlton N (2015) DNA barcoding and metabarcoding of standardized samples reveal patterns of marine benthic diversity. *Proceedings of the National Academy of Sciences* **112**, 2076-2081.
- Leray M, Meyer CP, Mills SC (2015) Metabarcoding dietary analysis of coral dwelling predatory fish demonstrates the minor contribution of coral mutualists to their highly partitioned, generalist diet. *PeerJ* **3**, e1047.
- Leray M, Yang JY, Meyer CP, *et al.* (2013) A new versatile primer set targeting a short fragment of the mitochondrial COI region for metabarcoding metazoan diversity: application for characterizing coral reef fish gut contents. *Frontiers in Zoology* **10**, 1-14.
- Luangsa-Ard J, Houbraken J, van Doorn T, *et al.* (2011) *Purpureocillium*, a new genus for the medically important *Paecilomyces lilacinus*. *FEMS Microbiology Letters* **321**, 141-149.
- Mahé F, Rognes T, Quince C, de Vargas C, Dunthorn M (2014) Swarm: robust and fast clustering method for amplicon-based studies. *PeerJ* **2**, e593.
- Mahé F, Rognes T, Quince C, de Vargas C, Dunthorn M (2015) Swarm v2: highly-scalable and high-resolution amplicon clustering. *PeerJ* **3**, e1420.
- Manter DK, Vivanco JM (2007) Use of the ITS primers, ITS1F and ITS4, to characterize fungal abundance and diversity in mixed-template samples by qPCR and length heterogeneity analysis. *Journal of Microbiological Methods* **71**, 7-14.
- Mauchline J (1971) The biology of *Neomysis integer* [Crustacea, Mysidacea]. *Journal of the Marine Biological Association of the United Kingdom* **51**, 347-354.
- Meadows PS (1964) Experiments on substrate selection by *Corophium Volutator* (Pallas): Depth selection and population density. *Journal of Experimental Biology* **41**, 677.
- Meadows PS, Reid A (1966) The behaviour of *Corophium volutator* (Crustacea: Amphipoda). *Journal of Zoology* **150**, 387-399.
- Mees J, Dewicke A, Hamerlynck O (1993) Seasonal composition and spatial distribution of hyperbenthic communities along estuarine gradients in the Westerschelde. *Netherlands Journal of Aquatic Ecology* **27**, 359-376.
- Moksnes PO (2002) The relative importance of habitat-specific settlement, predation and juvenile dispersal for distribution and abundance of young juvenile shore crabs *Carcinus maenas* L. *Journal of Experimental Marine Biology and Ecology* **271**, 41-73.
- Moksnes PO, Pihl L, van Montfrans J (1998) Predation on postlarvae and juveniles of the shore crab *Carcinus maenas*: Importance of shelter, size and cannibalism. *Marine Ecology Progress Series* **166**, 211-225.
- Molnar K, Ostoros G, Dunams-Morel D, Rosenthal BM (2012) Eimeria that infect fish are diverse and are related to, but distinct from, those that infect terrestrial vertebrates. *Infection, Genetics and Evolution* **12**, 1810-1815.

- Murray DC, Bunce M, Cannell BL, *et al.* (2011) DNA-based faecal dietary analysis: a comparison of qPCR and high throughput sequencing approaches. *PLoS One* **6**.
- Nidheesh T, Pal GK, Suresh PV (2015) Chitooligomers preparation by chitosanase produced under solid state fermentation using shrimp by-products as substrate. *Carbohydrate Polymers* **121**, 1-9.
- Nikolcheva LG, Bärlocher F (2004) Taxon-specific fungal primers reveal unexpectedly high diversity during leaf decomposition in a stream. *Mycological Progress* **3**, 41-49.
- Nordström M, Aarnio K, Bonsdorff E (2009) Temporal variability of a benthic food web: Patterns and processes in a low-diversity system. *Marine Ecology Progress Series* **378**, 13-26.
- Oh CW, Richard GH, Richard DMN (2001) Feeding ecology of the common shrimp *Crangon crangon* in Port Erin Bay, Isle of Man, Irish Sea. *Marine Ecology Progress Series* **214**, 211-223.
- Oksanen J, Guillaume Blanchet F, Kindt R, *et al.* (2016) Vegan: Community Ecology Package, pp. 631-637.
- Ólafsson EB, Persson L-E (1986) The interaction between *Nereis diversicolor* O.F. Müller and *Corophium volutator* Pallas as a structuring force in a shallow brackish sediment. *Journal of Experimental Marine Biology and Ecology* **103**, 103-117.
- Olmos-Perez L, Roura A, Pierce GJ, Boyer S, Gonzalez AF (2017) Diet composition and variability of wild *Octopus vulgaris* and *Alloteuthis media* (Cephalopoda) Paralarvae: a metagenomic approach. *Frontiers in Physiology* **8**, 321.
- Pallas A, Garcia-Calvo B, Corgos A, Bernardez C, Freire J (2006) Distribution and habitat use patterns of benthic decapod crustaceans in shallow waters: a comparative approach. *Marine Ecology Progress Series* **324**, 173-184.
- Peterson WT, Ausubel SJ (1984) Diets and selective feeding by larvae of Atlantic mackerel *Scomber scombrus* on zooplankton *Marine Ecology Progress Series* **17**, 65-75.
- Pihl L (1985) Food selection and consumption of mobile epibenthic fauna in shallow marine areas. *Marine Ecology Progress Series* **22**, 169-179.
- Pihl L, Rosenberg R (1984) Food selection and consumption of the shrimp *Crangon crangon* in some shallow marine areas in western Sweden. *Marine Ecology Progress Series* **15**, 159-168.
- Pinn EH, Ansell AD (1993) The effect of particle size on the burying ability of the brown shrimp *Crangon crangon*. *Journal of the Marine Biological Association of the United Kingdom* **73**, 365-377.
- Pinol J, Mir G, Gomez-Polo P, Agusti N (2015) Universal and blocking primer mismatches limit the use of high-throughput DNA sequencing for the quantitative metabarcoding of arthropods. *Molecular Ecology Resources* **15**, 819-830.
- Pinol J, San Andres V, Clare EL, Mir G, Symondson WO (2014) A pragmatic approach to the analysis of diets of generalist predators: the use of next-generation sequencing with no blocking probes. *Molecular Ecology Resources* **14**, 18-26.
- Plagmann J (1939) Ernährungsbiologie der Garnele (*Crangon vulgaris* Fabr.). *Helgoländer wissenschaftliche Meeresuntersuchungen* **2**, 113-162.
- Pompanon F, Deagle BE, Symondson WOC, *et al.* (2012) Who is eating what: diet assessment using next generation sequencing. *Molecular Ecology* **21**.
- Price KS (1962) Biology of the sand shrimp, *Crangon septemspinosa*, in the shore zone of the Delaware Bay Region. *Chesapeake Science* **3**, 244-255.
- Raffaelli D, Conacher A, McLachlan H, Emes C (1989) The role of epibenthic crustacean predators in an estuarine food web. *Estuarine, Coastal and Shelf Science* **28**, 149-160.
- Ratnasingham S, Hebert PDN (2007) BOLD: The Barcode of Life Data System (<http://www.barcodinglife.org>). *Molecular Ecology Notes* **7**, 355-364.
- Ray JL, Althammer J, Skaar KS, *et al.* (2016) Metabarcoding and metabolome analyses of copepod grazing reveal feeding preference and linkage to metabolite classes in dynamic microbial plankton communities. *Molecular Ecology* **25**, 5585-5602.

- Redou V, Navarri M, Meslet-Cladiere L, Barbier G, Burgaud G (2015) Species richness and adaptation of marine fungi from deep-subseafloor sediments. *Applied and Environmental Microbiology* **81**, 3571-3583.
- Rodrigues AM, Quintino V, Sampaio L, Freitas R, Neves R (2011) Benthic biodiversity patterns in Ria de Aveiro, Western Portugal: Environmental-biological relationships. *Estuarine, Coastal and Shelf Science* **95**, 338-348.
- Rognes T, Flouri T, Nichols B, Quince C, Mahe F (2016) VSEARCH: a versatile open source tool for metagenomics. *PeerJ* **4**, e2584.
- Rueckert S, Simdyanov TG, Aleoshin VV, Leander BS (2011) Identification of a divergent environmental DNA sequence clade using the phylogeny of gregarine parasites (apicomplexa) from crustacean hosts. *PLoS One* **6**, e18163.
- Sakaguchi SO, Shimamura S, Shimizu Y, *et al.* (2017) Comparison of morphological and DNA-based techniques for stomach content analyses in juvenile chum salmon *Oncorhynchus keta*: a case study on diet richness of juvenile fishes. *Fisheries Science* **83**, 47-56.
- Seikai T, Kinoshita I, Tanaka M (1993) Predation by crangonid shrimp on juvenile japanese flounder under laboratory conditions. *Nippon Suisan Gakkaishi* **59**, 321-326.
- Singh S, Pandey RK, Goswami BK (2013) Bio-control activity of *Purpureocillium lilacinum* strains in managing root-knot disease of tomato caused by *Meloidogyne incognita*. *Biocontrol Science and Technology* **23**, 1469-1489.
- Smith AL, Hamilton KM, Hirschle L, *et al.* (2013) Characterization and molecular epidemiology of a fungal infection of edible crabs (*Cancer pagurus*) and interaction of the fungus with the dinoflagellate parasite *Hematodinium*. *Applied and Environmental Microbiology* **79**, 783-793.
- Soininen EM, Zinger L, Gielly L, *et al.* (2013) Shedding new light on the diet of Norwegian lemmings: DNA metabarcoding of stomach content. *Polar Biology* **36**, 1069-1076.
- Stentiford GD, Shields JD (2005) A review of the parasitic dinoflagellates *Hematodinium* species and *Hematodinium*-like infections in marine crustaceans. *Diseases of Aquatic Organisms* **66**, 47-70.
- Strauss RE (1979) Reliability estimates for Ivlev's Electivity Index, the Forage Ratio, and a Proposed Linear Index of Food Selection. *Transactions of the American Fisheries Society* **108**, 344-352.
- Symondson WOC (2002) Molecular identification of prey in predator diets. *Molecular Ecology* **11**.
- Taberlet P, Coissac E, Pompanon F, Brochmann C, Willerslev E (2012a) Towards next-generation biodiversity assessment using DNA metabarcoding. *Molecular Ecology* **21**, 2045-2050.
- Taberlet P, Prud'Homme SM, Campione E, *et al.* (2012b) Soil sampling and isolation of extracellular DNA from large amount of starting material suitable for metabarcoding studies. *Molecular Ecology* **21**, 1816-1820.
- Tang CQ, Leasi F, Obertegger U, *et al.* (2012) The widely used small subunit 18S rDNA molecule greatly underestimates true diversity in biodiversity surveys of the meiofauna. *Proceedings of the National Academy of Sciences* **109**, 16208-16212.
- Thomas AC, Deagle BE, Eveson JP, Harsch CH, Trites AW (2016) Quantitative DNA metabarcoding: improved estimates of species proportional biomass using correction factors derived from control material. *Molecular Ecology Resources* **16**, 714-726.
- Tiews K (1970) Synopsis of biological data on the common shrimp *Crangon crangon* Linnaeus 1758. *FAO Fisheries Reports* **4**, 1167-1224.
- Turner CR, Uy KL, Everhart RC (2015) Fish environmental DNA is more concentrated in aquatic sediments than surface water. *Biological Conservation* **183**, 93-102.
- Ushio M, Murakami H, Masuda R, *et al.* (2018) Quantitative monitoring of multispecies fish environmental DNA using high-throughput sequencing. *Metabarcoding and Metagenomics* **2**, e23297.
- Vamos EE, Elbrecht V, Leese F (2017) Short COI markers for freshwater macroinvertebrate metabarcoding. *Metabarcoding and Metagenomics* **1**, e14625.

- van der Veer HW, Bergman MJ (1987) Predation by crustaceans on a newly settled 0-group plaice *Pleuronectes platessa* population in the western Wadden Sea. *Marine Ecology Progress Series* **35**, 203-215.
- van der Veer HW, Feller RJ, Weber A, Witte J (1998) Importance of predation by crustaceans upon bivalve spat in the intertidal zone of the Dutch Wadden Sea as revealed by immunological assays of gut contents. *Journal of Experimental Marine Biology and Ecology* **231**, 139-157.
- Walter U, Becker PH (1997) Occurrence and consumption of seabirds scavenging on shrimp trawler discards in the Wadden Sea. *ICES Journal of Marine Science: Journal du Conseil* **54**, 684-694.
- Wangensteen O, Turon X (2017) Metabarcoding techniques for assessing biodiversity of marine animal forests. In: *Marine animal forests. The ecology of benthic biodiversity hotspots* (eds. Rossi S, Bramanti L, Gori A, Orejas C), pp. 445-473. Springer International Publishing. (Print ISBN: 978-3-319-21011-7).
- Wangensteen OS, Palacín C, Guardiola M, Turon X (2018) DNA metabarcoding of littoral hard-bottom communities: high diversity and database gaps revealed by two molecular markers. *PeerJ* **6**, e4705.
- Wilcox JR, Jeffries HP (1974) Feeding habits of the sand shrimp *Crangon septemspinosa*. *Biological Bulletin* **146**, 424-434.
- Yue Y, Yu H, Li R, *et al.* (2015) Exploring the antibacterial and antifungal potential of jellyfish-associated marine fungi by cultivation-dependent approaches. *PLoS One* **10**, e0144394.

813 **Data Accessibility**

814 The dataset, including sequences, taxonomic assignment and abundances for all MOTUs in every
815 sample, have been deposited in Dryad (doi:10.5061/dryad.sk2155m). Custom R scripts are publicly
816 available from <https://github.com/Andjin/Crangon-diet-analysis> and
817 <https://github.com/metabarpark> for scripts related to the bioinformatics pipeline.

818


819 **Author Contributions**

820 A.S. participated in the study design, carried out the field, laboratory and molecular work, performed
821 the data and statistical analyses, and wrote the manuscript; O.S.W. participated in the molecular work,
822 designed the bioinformatics pipeline and assisted in the statistical analyses; C.B. participated in the
823 study design; J.C. assisted in the collection of the samples; S.M. conceived and designed the study. All
824 authors provided critical comments on the manuscript and gave final approval for publication.

825

826

827



831 Figure 1. Overview of sample locations, illustrating (A) the overall western European scale; (B) the
832 Dutch estuaries, Western Scheldt (WS) and Eastern Scheldt (ES); (C) the British estuaries, Mersey (Me)
833 and Kent (Ke); the Aveiro (D) and Minho (E) estuaries in Portugal. Small dots within estuaries represent
834 individual collection points for shrimp and sediment samples. *Site removed prior to molecular
835 analysis. Source map: OpenStreetMap.




Figure 2. Relative abundances of MOTUs detected in *Crangon crangon* stomach samples by COI metabarcoding, after removal of *C. crangon* and *P. lilacinum* reads. Each bar represents one sample. Countries are shown on top of the graph, estuaries below and boxes contain the individual sites. The number on top of each sample represents the number of diet-related COI reads. The category other is comprised of MOTUs with < 1.0 % COI reads.

843 Table 1. Trophic significance of *Crangon crangon* prey items. MOTUs shown (≥ 0.5 % mean relative abundance in the stomach samples) are assigned to the
844 family level or lower.

Phylum	Order	Family	Species	Best identity	Presence (%)	Mean (\pm SE) abundance (%)	Mean (\pm SE) selectivity (D)	Trophic significance	Literature	Source
Annelida	Phyllodocida	Nereididae	<i>Hediste diversicolor</i>	0.98	30.8	7.1 \pm 4.5	0.9 \pm 0.0	Medium	Sto+	Lloyd and Yonge (1947); Pihl and Rosenberg (1984)
	Spionida	Spionidae	<i>Scolecipis foliosa</i>	1.00	5.8	0.7 \pm 0.6	1.0 \pm 0.0	Low	Exp ² Sto- ²	Van Tomme et al. (2014); Ansell et al. (1999)
	Terebellida	Terebellidae	<i>Pista cristata</i>	0.99	21.2	4.5 \pm 2.6	0.6 \pm 0.2	Medium	Sto- ²	Ansell et al. (1999)
Arthropoda	Amphipoda	Corophiidae	<i>Corophium volutator</i>	0.99	28.8	13.8\pm5.9	1.0\pm0.0	High	Sto+	Pihl and Rosenberg (1984); Evans (1984)
		Pontoporeiidae	<i>Bathyporeia sarsi</i>	1.00	7.7	3.5 \pm 2.5	1.0 \pm 0.0	Medium	Exp	Van Tomme et al. (2014)
		Talitridae	<i>Talitrus saltator</i>	0.99	1.9	0.8 \pm 0.8	1.0 \pm 0.0	Low		
		Calanoida	Centropagidae	1.00	5.8	0.7 \pm 0.7	1.0 \pm 0.0	Low		
	Decapoda	Carcinidae	<i>Carcinus maenas</i>	1.00	55.8	8.0\pm3.3	1.0\pm0.0	High	Exp ¹ Sto-	Moksnes et al. (1998); Raffaelli et al. (1989); Pihl and Rosenberg (1984)
		Porcellanidae	<i>Pisidia longicornis</i>	1.00	9.6	1.7 \pm 1.5	1.0 \pm 0.0	Medium		
		Chironomidae	Unassigned	0.88	53.8	0.9 \pm 0.4	0.8 \pm 0.1	Medium	CA	Nordström et al. (2009)
	Mysida	Chironomidae	<i>Chironomus salinaris</i>	1.00	1.9	1.3 \pm 1.3	1.0 \pm 0.0	Low		
		Mysidae	<i>Mesopodopsis slabberi</i>	0.99	3.8	0.6 \pm 0.6	1.0 \pm 0.0	Low		
		Mysidae	<i>Neomysis integer</i>	0.98	13.5	2.7 \pm 2.1	1.0 \pm 0.0	Medium	Sto-	Raffaelli et al. (1989)
		Mysidae	<i>Schistomysis ornata</i>	0.98	1.9	1.6 \pm 1.6	1.0 \pm 0.0	Low	Sto+ ²	Oh et al. (2001)
	Sessilia	Austrobalanidae	<i>Austrominius modestus</i>	1.00	17.3	2.0 \pm 1.5	1.0 \pm 0.0	Medium		
Chordata	Atheriniformes	Atherinidae	<i>Atherina presbyter</i>	1.00	1.9	0.6 \pm 0.6	1.0 \pm 0.0	Low		
	Clupeiformes	Clupeidae	<i>Sardina pilchardus</i>	1.00	3.8	0.6 \pm 0.6	1.0 \pm 0.0	Low		
	Pleuronectiformes	Scophthalmidae	<i>Zeugopterus punctatus</i>	0.99	3.8	0.8 \pm 0.6	1.0 \pm 0.0	Low		
	Scombriformes	Scombridae	<i>Scomber scombrus</i>	1.00	1.9	1.0 \pm 1.0	1.0 \pm 0.0	Low		
	Spariformes	Sparidae	<i>Spondyliosoma cantharus</i>	1.00	7.7	0.6 \pm 0.4	1.0 \pm 0.0	Low		
Cnidaria	Actiniaria	Actiniidae	<i>Anthopleura elegantissima</i>	0.99	3.8	0.6 \pm 0.6	1.0 \pm 0.0	Low		

845 Exp: Experimental study; Sto+: Major contributor based on stomach analysis; Sto-: Minor contributor based on stomach analysis; CA: Contribution assumed by source; 1: Larvae; 2: Related taxa (same family). In bold: High trophic
846 significant taxa. Best identity: Alignment score of the best match in the reference database.




Figure 3. Relative abundances of MOTUs detected in sediment samples by COI metabarcoding. Each bar represents one sample. Countries are shown on top of the graph, estuaries below. The number on top of each sample represents the number of COI reads. The category other is comprised of MOTUs with < 1.0 % COI reads.





Figure 4. Phyla detected in sediment and *Crangon crangon* stomach samples by COI metabarcoding. (A) Total number of MOTU detected per phylum in sediment, full stomachs and visually empty stomachs. (B) Mean relative read abundance of phyla detected in sediment and *C. crangon* full stomach samples, after removal of *C. crangon* and *P. lilacinum* reads. (C) Phylum trophic significance based on presence (%), mean relative abundance (%) in full stomach samples and Jacobs' selectivity index. Stomach samples consisted of a pool of up to 8 stomachs. The category "other phyla" (represented in white) contains phyla with < 1% COI reads in both the sediment and full stomach samples.

862



863

864 Figure 5. Multidimensional scaling analysis of MOTUs detected in sediment (dots) and *Crangon*
 865 *crangon* stomach samples (triangles), based on square-root transformed Bray-Curtis dissimilarities.
 866 75% confidence ellipses are shown per sample type. WS = Western Scheldt; ES = Eastern Scheldt.

867




Figure 6. Multivariate analysis of *Crangon crangon* diet in six estuaries determined by COI metabarcoding based on MOTUs (N = 20) over all stomach samples (N = 53). (A) Mean relative read abundance of each MOTU per estuary based on DNA extracted from sediment and stomach samples (after removal of *C. crangon* and *P. lilacinum* reads). MOTUs are identified for $\geq 1.0\%$ average read abundance in the stomach samples, otherwise are referred as “Other” (B) Non-metric multidimensional scaling (nMDS) analysis based on Bray-Curtis dissimilarities of square-root transformed relative abundances in *C. crangon* stomach samples. Each dot represents one pooled stomach sample, estuaries are identified by colours (see below) and ellipses show 75% confidence intervals. (C) Canonical correspondence analysis (CCA) of square-root transformed relative read abundances in relation to salinity, total organic matter (TOM) and median grain size. Reads were averaged per site (displayed as dots) and estuaries are identified by colour (see below). Red crosses represent the MOTU scores and numbers refer to the MOTU names given in panel A.




Figure 7. MOTU accumulation curves showing MOTU richness (based on all MOTUs detected) in *Crangon crangon* pooled stomach samples in several European estuaries. Each sample has been rarefied to 1000 reads prior to the construction of the accumulation curves. Stomach samples consisted of a pool of up to 8 stomachs.




Figure 8. Schematic representation of the most important food items of adult *Crangon crangon* and their probable method of capture/ingestion. Line thickness represents trophic significance: high (bold); medium (thin); low (dashed). Numbers identify prey categories: annelids (1); decapod larvae/instars (2); fish 0-year-juveniles (3); 4 mysids (4); (pico) phytoplankton (5); fish carcasses (6); decapod carcasses (7); amphipods (8); chironomid, mollusc and barnacle larvae (9); meiofauna (10). Letters define method of ingestion: Secondary predation (SP); ambush predation (AP); gulping predation (GP); passive ingestion (PI); scavenging (SC). Images not to scale.