



Mitochondrial DNA Part B

Resources

ISSN: (Print) 2380-2359 (Online) Journal homepage: https://www.tandfonline.com/loi/tmdn20

Complete mitochondrial DNA sequence of the tropical hornet *Vespa affinis* (Insecta, Hymenoptera)

Hisashi Okuyama, Stephen J. Martin & Jun-Ichi Takahashi

To cite this article: Hisashi Okuyama, Stephen J. Martin & Jun-Ichi Takahashi (2017) Complete mitochondrial DNA sequence of the tropical hornet *Vespa affinis* (Insecta, Hymenoptera), Mitochondrial DNA Part B, 2:2, 776-777, DOI: <u>10.1080/23802359.2017.1398622</u>

To link to this article: <u>https://doi.org/10.1080/23802359.2017.1398622</u>

© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 07 Nov 2017.

|--|

Submit your article to this journal \square

Article views: 163



View Crossmark data 🗹

👖 Citing articles: 2 View citing articles 🗹

MITOGENOME ANNOUNCEMENT

OPEN ACCESS Check for updates

Taylor & Francis

Taylor & Francis Group

Complete mitochondrial DNA sequence of the tropical hornet *Vespa affinis* (Insecta, Hymenoptera)

Hisashi Okuyama^a, Stephen J. Martin^b and Jun-Ichi Takahashi^a

^aDepartment of Life sciences, Kyoto Sangyo University, Kyoto, Japan; ^bSchool of Environment & Life Sciences, University of Salford, Manchester, UK

ABSTRACT

We analyzed the complete mitochondrial genome of the Asian tropical hornet *Vespa affinis* from Ishigaki Island, Japan. It consisted of a circular molecule with 19,109 bp, which is larger to other hornet species e.g. *V. velutina*. We predicted the genome contained 13 protein-coding, 22 tRNA, and two rRNA genes, along with one A + T-rich control region. The repetitive sequences were confirmed at multiple positions in the non-coding genes. The initiation codons ATA was found in one, ATG in seven, and ATT in five genes, while the termination codons TAA and TAG were observed 11 and two genes, respectively. The average AT content of the genome was 78.4%.

ARTICLE HISTORY

Received 22 October 2017 Accepted 26 October 2017

KEYWORDS

Hornet; Illumina sequencing; repetitive sequence; monogynous; polygynous

There are 22 hornet (genus Vespa) species (Archer 2012; Perrard et al. 2013) and these have become an important model group for research into social systems such as worker altruistic behaviours, reproductive conflict, evolution of the queen number and development of sociality (Matsuura and Yamane 1990; Foster and Ratnieks 2001). The tropical hornet Vespa affinis is distributed throughout the subtropical and tropical regions of Asia (Archer 2012). Normally the V. affinis colony is founded by a single queen, although some nests contain multiple queens in tropical regions (Matsuura and Yamane 1990; Martin 1995a, 1995b), an unusual trait among the hornets (Martin et al. 2009). To date, there has been limited analysis of mitochondrial DNA from hornets (Takeuchi et al. 2017), so hampering a detailed knowledge about their population genetic structure, intraspecific phylogeny relationships, and genetic diversity. Therefore, to help resolve this issue, we analyzed the complete mitochondrial genome of the hornet V. affinis.

A single adult worker was collected from a colony on Ishigaki Island, Okinawa Prefecture, in the sub-tropical region of Japan (the specimen was stored in the National Museum of Nature and Science, Japan, accession number: NSMT-I-HYM 75315). Genomic DNA was isolated and sequenced using Illumina's Next Seq 500 (Illumina Inc, San Diego, CA). The 1,462,186 reads were assembled and annotated using the MITOS web server (Bernt et al. 2013), MEGA6 (Tamura et al. 2013), and GNETYX v.10 (Genetyx Corporation, Tokyo, Japan). The phylogenetic analysis was performed using TREEFINDER (March 2011) (Jobb et al. 2004) based on the nucleotide sequences of the 13 protein-coding genes.

The V. affinis mitochondrial genome forms a 19,109 bplong closed loop (accession number AP018371). Although it is \sim 2000 bp longer than the *Vespa* genomes it has a very similar genomic organization, since it is composed of 13 protein-coding, 22 putative tRNA, and two rRNA genes, as well as an A+T-rich control region. The average AT content of the 13 protein-coding genes was 78.4%, similar to the 82% found in V. velutina (Takahashi et al. 2017). The V. affinis genome, was predicted to have nine protein-coding and 14 tRNA genes on the heavy strand and four protein-coding, eight tRNA, and two rRNA genes on the light strand. The genes ND4 and ND4L shared seven nucleotides. Of the 13 proteincoding genes, the initiation codon ATA was found in one, ATG in seven, and ATT in five genes, while the termination codon TAA and TAG were observed in the protein-coding genes ND4 and Cytb, respectively.

Phylogenetic analysis using the 13 mitochondrial proteincoding genes from 11 closely related taxa of Vespidae (Cameron et al. 2008; Chen et al. 2016; Song et al. 2016; Wei et al. 2016; Zhou et al. 2016; Fan et al. 2017; Haddad et al. 2017; Kim et al. 2017a, 2017b; Takahashi et al. 2017) was shown to be similar to the result by the combined analysis based on 45 morphological characters and six genes (Perrard et al. 2013) (Figure 1). Although more high resolution sequence data from more species may help resolve these differences. Finally, non-coding repetitive sequences in the mitochondrial DNA of *V. affinis* were confirmed at multiple positions between the 13 protein-coding genes and these regions will help to develop a suite of specific primers for the estimation of genetic relationship and genetic diversity.

CONTACT Jun-ichi Takahashi 🖾 jit@cc.kyoto-su.ac.jp 🗗 Department of Life Sciences, Kyoto Sangyo University, Kyoto, Japan

© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Figure 1. Phylogenetic relationships (maximum likelihood) of the Vespidae based on the nucleotide sequence of 13 protein-coding genes of the mitochondrial genome. The numbers at the nodes indicate bootstrap support inferred from 1000 bootstrap replicates. The sequence of Abispa ephippium was used as outgroup. Alphanumeric terms indicate the GenBank accession numbers.

Acknowledgements

The authors thank Mr. Unebi Enami for helping our research.

Disclosure statement

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of this article.

Funding

This study was supported by JSPS KAKENHI Grant no. 17K07575.

References

- Archer ME. 2012. Vespine wasps of the world: behavior, ecology & taxonomy of the Vespinae. Manchester: Siri Scientific Press.
- Bernt M, Donath A, Juhling F, Externbrink F, Florentz C, Fritzsch G, Putz J, Middendorf M, Stadler PF. 2013. MITOS: improved de novo Metazoan Mitochondrial Genome Annotation. Mol Phylogenet Evol. 69:313–319.
- Cameron SL, Dowton M, Castro LR, Ruberu K, Whiting MF, Austin AD, Diement K, Stevens J. 2008. Mitochondrial genome organization and phylogeny of two vespid wasps. Genome. 51:800–808.
- Chen PY, Wei SJ, Liu JX. 2016. The mitochondrial genome of the *Vespa* mandarinia Smith (Hymenoptera: Vespidae: Vespinae) and a phylogenetic analysis of the Vespoidea. Mitochondrial DNA Part A. 27:4414–4415.
- Fan XL, Gong YJ, Chen PY, Tan QQ, Tan JL, Wet SJ. 2017. Next-generation sequencing of the mitochondrial genome of *Dolichovespula panda* (Hymenoptera: Vespidae) with a phylogenetic analysis of Vespidae. J Asia Pac Entomol. 20:971–976.
- Foster KR, Ratnieks FLW. 2001. Paternity, reproduction and conflict in vespine wasps: a model system for testing kin selection predictions. Behav Ecol Sociobiol. 50:1–8.
- Haddad NJ, Al-Nakeeb K, Petersen B, Dalén L, Blom N, Sicheritz-Pontén T.
 2017. Complete mitochondrial genome of the Oriental Hornet, *Vespa* orientalis F. (Hymenoptera: Vespidae). Mitochondrial DNA Part B. 2:139–140.
- Jobb G, von Haeseler A, Strimmer K. 2004. TREEFINDER: a powerful graphical analysis environment for molecular phylogenetics. BMC Evol Biol. 4:18.

- Kim JS, Jeong JS, Kim I. 2017a. Complete mitochondrial genome of the yellow-legged Asian hornet, *Vespa velutina nigrithorax* (Hymenoptera: Vespidae). Mitochondrial DNA Part B. 2:82–84.
- Kim JS, Jeong JS, Jeong SY, Kim MJ, Kim I. 2017b. Complete mitochondrial genome of the black-tailed hornet, *Vespa ducalis* (Hymenoptera: Vespidae): genomic comparisons in Vespoidea. Entomol Res. 47:129–136.
- Martin SJ. 1995a. Colony development of the hornet *Vespa affinis* (Hymenoptera: Vespidae). Jpn J Entomol. 63:861–876.
- Martin SJ. 1995b. Comparison of tropical and sub-tropical colonies of the hornet *Vespa affinis*. Asian J Tropic Biol. 2:13–19.
- Martin SJ, Takahashi J, Katada S. 2009. Queen condition, mating frequency, queen loss, and levels of worker reproduction in the hornets *Vespa affinis* and *V. simillima*. Ecol Entomol. 34:43–49.
- Matsuura M, Yamane S. 1990. Biology of the Vespine Wasps. Berlin: Springer.
- Perrard A, Pickett KM, Villemant C, Kojima J, Carpenter J. 2013. Phylogeny of hornets: a total evidence approach (Hymenoptera, Vespidae, Vespinae, Vespa). J Hymenoptera Res. 32:1–15.
- Song SN, Chen PY, Wei SJ, Chen XX. 2016. The mitochondrial genome of *Polistes jokahamae* and a phylogenetic analysis of the Vespoidea (Insecta: Hymenoptera). Mitochondrial DNA Part A. 27:2783–2784.
- Takahashi R, Okuyama H, Kiyoshi K, Takahashi J. 2017. Complete mitochondrial DNA sequence of the invasive hornet *Vespa velutina* (Insecta, Hymenoptera) found in Japan. Mitochondrial DNA Part B. 2:143–144.
- Takeuchi T, Takahashi R, Kiyoshi T, Nakamura M, Minoshima Y, Takahashi J. 2017. The origin and genetic diversity of the yellow-legged hornet, *Vespa velutina* introduced in Japan. Insect Soc. 64:313–320.
- Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013. MEGA6: Molecular evolutionary genetics analysis version 6.0. Mol Biol Evol. 30:2725–2729.
- Wei SJ, Niu FF, Tan JL. 2016. The mitochondrial genome of the Vespa bicolor Fabricius (Hymenoptera: Vespidae: Vespinae). Mitochondrial DNA Part A. 27:875–876.
- Zhou Y, Hu YL, Xu ZF. 2016. The mitochondrial genome of the German wasp Vespula germanica (Fabricius, 1793) (Hymenoptera: Vespoidea: Vespidae). Mitochondrial DNA Part A. 27:2917–2918.