

# Coupled spatiotemporal waves: new paradigms in vector soliton physics


J. T. Ashley, J. M. Christian, and G. S. McDonald

*Materials & Physics Research Centre, University of Salford, U.K.*

Email: [j.t.ashley@edu.salford.ac.uk](mailto:j.t.ashley@edu.salford.ac.uk)

Keywords: vector solitons, spatiotemporal dispersion, *energy theme*

In this presentation, we propose a novel spatiotemporal generalization of Menyuk's classic equations [1] describing the propagation of two nonlinearly-coupled waves in a dispersive optical system (such as a fibre or planar waveguide). Our approach is similar in spirit to that recently introduced by Christian *et al.* [2] in a single-component context: both analyses are firmly rooted in special relativity-inspired frame-of-reference considerations, and are concerned with the universal properties of waves in fully second-order systems (where evolution is allowed to venture beyond the realms of the ubiquitous slowly-varying envelope approximation). Research highlights include the complete solution of the modulational instability (MI) problem for vectorized spatiotemporal continuous-wave solutions (the characteristic equation turns out to be an eighth-degree polynomial, rather than a quartic [1]). Predictions of the most-unstable frequency in the system, obtained from consultation of the MI spectrum, are in full agreement with extensive numerical calculations [see Figs. 1(a) & 1(b)]. We have also derived exact bright-bright solitons and investigated some of their key characteristics (e.g., robustness and the asymptotic recovery of Menyuk-type solutions). Future work will address spatiotemporal vector dark solitons [3,4] [see Fig. 1(c)].



**Figure 1.** (a) Typical long-wave MI spectrum when both waves experience anomalous temporal dispersion. (b) Extensive simulations have confirmed the prediction for the most-unstable frequency, denoted by  $\Omega_{P0}$  [related to the dominant peak in part (a)]. (c) Instability of a conventional dark-dark type of initial condition used for the spatiotemporal governing equation.

## References

- [1] C. R. Menyuk, IEEE J. Quantum Electron. **QE-23**, 174 (1987).
- [2] J. M. Christian, G. S. McDonald, T. F. Hodgkinson, and P. Chamorro-Posada, Phys. Rev. Lett. **108**, 034101 (2012).
- [3] V. V. Afanasyev, Y. S. Kivshar, V. V. Konotop, and V. N. Serkin, Opt. Lett. **14**, 805 (1989).
- [4] Y. S. Kivshar and S. K. Turitsyn, Opt. Lett. **18**, 337 (1993).