

1 **Occurrence of Deformed wing virus variants in the stingless *Melipona subnitida* and honey**
2 ***Apis mellifera* bee populations in North Eastern Brazil**

3
4 Flaviane S. de Souza^{1,2}, Jessica L. Kevill¹, Carlos A. L. de Carvalho², Stephen J. Martin^{1*}

5
6 ¹*School of Environment and Life Sciences, The University of Salford, Manchester M5 4WT, UK.*

7
8 ²*Universidade Federal do Recôncavo da Bahia, Rua Rui Barbosa 710, 44380-000, Cruz das*
9 *Almas, Bahia, Brazil*

10
11 *Corresponding author: s.j.martin@salford.ac.uk

12
13 Key Words: spill-over, Varroa, viral variants

14
15 Abbreviations DWV, Deformed wing virus

16
17 **Abstract**

18 Deformed wing virus (DWV) is now a global insect pathogen. Brazilian stingless bees are a
19 diverse group often managed in close proximity to honey bees. We investigated the prevalence
20 and load of DWV in 33 stingless bees (*Melipona subnitida*) and 12 honey bees (*Apis mellifera*)
21 colonies from NE Brazil. DWV was detected in all colonies with the A and C-variants dominating
22 *M. subnitida* and A-variant in *A. mellifera*. Viral loads were 8.83E+07 and 7.19E+07 in *M.*
23 *subnitida* and *A. mellifera*, respectively. On Fernando de Noronha island DWV is low (<1E+03)
24 in honey bees, but we detected high loads (1.6E+08) in nine island *M. subnitida* colonies,
25 indicating no viral spill-over of DWV has occurred during the past 34 years. Furthermore, the
26 ubiquitous presence of the DWV-C variant in *M. subnitida* colonies, and rarity in *A. mellifera*,
27 may suggest limited viral exchange between these two species.

36 INTRODUCTION

37 The stingless bees (Apidae: *Meliponini*) are the most diverse group of eusocial bees,
38 comprising of more than 400 species contained within 60 genera [1]. The majority of species
39 occur in the Neo-tropics with colonies typically containing 200-700 adults and a perennial life-
40 cycle [2]. Many species, particularly the large *Melipona* species have a long association with
41 humans that harvest their highly prized honey [3], but they are also responsible for pollinating
42 40-90% of the native flora in some regions of Brazil [4]. Relative to the honey bees (*Apis* spp),
43 very little is known about the pests and pathogens of stingless bees despite their importance.

44 Brazil has a long history of managing honey bees (*Apis mellifera*) originally imported
45 from Europe, but in 1957, 26 colonies of imported African *A. m. scutellata* escaped quarantine
46 and spread throughout Brazil, hybridising with existing honeybees to form the Africanised honey
47 bee [5]. However, when in 1971 the parasitic Varroa (*Varroa destructor*) mite arrived in Brazil,
48 the Africanised honey bees were naturally tolerant to the mite, whereas, the European honeybees
49 suffered large scale losses. These losses are caused by a viral pathogen called Deformed Wing
50 Virus (DWV) that is transmitted by the Varroa mite [6].

51 Although Varroa can only survive on honey bees, [7] showed that raised DWV levels in
52 the honey bee population, initiated by the mite, has resulted in viral spill-over into other species
53 of bees and wasps. This may explain why DWV has been detected in a wide range of non-*Apis*
54 insects [8-11] and has even been detected in pollen [12]. The impact of DWV on these hosts
55 remains unknown [13], although there is growing concern [11, 14-16].

56 In Brazil, the Africanised honey bee, Varroa mite and DWV have been present for
57 decades so there have been ample opportunities for cross-species infections to occur, especially
58 since both honey bees and stingless bees are often managed in close proximity, i.e. in nearby
59 apiaries. Therefore, the aim of this study was to evaluate both the prevalence and viral load of
60 the three described DWV master-variants (A, B and C) across a population of stingless bee
61 (*Melipona subnitida*) and Africanised honey bees from North-Eastern Brazil. The stingless bee
62 *M. subnitida* is a swarm founding species, brood development takes around 40 days, and workers
63 survive for a few months. This species is endemic to the dryland-shrub forest 'Caatinga biome'
64 found in NE Brazil and is the typical stingless bee maintained by beekeepers throughout the
65 region. This Meliponiculture helps towards the conservation of local biodiversity, as well as
66 provide extra income to the beekeepers [3].

70 RESULTS

71 Prevalence of DWV

72 We detected DWV in every *M. subnitida* and *A. mellifera* colony. Negative controls
73 indicated no contamination had occurred in any of the runs. Furthermore, the housekeeping gene
74 indicated all samples contain intact RNA (Fig. 1). The average Ct values indicated more β -actin
75 in the *A. mellifera* samples ($19.7\text{Ct} \pm 1.91$ S.D.) relative to the *M. subnitida* samples ($23.5\text{Ct} \pm$
76 0.70 S.D.).

77
78 **DWV viral loads**, The A and C master-variants were detected in the *M. subnitida* population only
79 (Fig. 2). The DWV-A variant was dominant in 78% of the colonies (Fig. 2) with the C-variant
80 dominating the remaining 22%. Whereas, 92 % of honey bee colonies were dominated by the A-
81 variant and only one colony (8%) was dominated by the C-variant. The DWV-B variant was
82 quantifiable in a single *A. mellifera* colony (Table 1) whilst three others tested positive below the
83 quantifiable limit of the qPCR assay but had visible bands when visualised on a gel (Mossoro,
84 Garanhus and Cruz das Almas). The total DWV viral load detected in both species of bee averaged
85 $8.8\text{E}+07$ and $7.2\text{E}+07$ in *M. subnitida* and *A. mellifera* respectively. On the remote Fernando de
86 Noronha island, the *M. subnitida* colonies were dominated by the A-variant, and the C-variant was
87 widespread. However, the viral load was an order of magnitude higher on the island ($1.6\text{E}+08$)
88 relative to the mainland ($3.6\text{E}+07$).

89

90 DISCUSSION

91 This study provides the first report of DWV in *Meliponini* stingless bees, since DWV was
92 not detected previously in *Melipona quadrifasciata* and *M. torrida* [17], although the DWV-A
93 variant was detected in Argentinian stingless bees (*Tetragonisca fiebrigi*). Furthermore, *M.*
94 *scutellaris* tested negative for six bee-associated viruses including DWV, but did test positive for
95 the honey bee associated acute bee paralysis virus [18]. The high prevalence of DWV in *A.*
96 *mellifera* was expected since DWV is consistently the most prevalent viral pathogen of European
97 and Africanised honey bees [19].

98 The dominance of the DWV-A variant found in this study reflects the situation found in
99 honey bees in the USA in 2010 [20]. Although the B-variant is replacing the A-variant in the USA
100 [20] and appears common in Europe [21], it was only detected in any quantity in a single
101 Africanised colony (Fig. 2). This is despite the likely long-term infection of both stingless and
102 honey bees in Brazil. The rarely detected C-variant [20, 22] was present in almost all the *M.*
103 *subnitida* colonies.

104 Interestingly on the remote island of Fernando de Noronha where both *M. subnitida* and
105 *A. mellifera* have been maintained in close proximity over the past 34 years, the DWV-A variant
106 dominated all nine colonies with a mean viral load of 1.6E+08. Whereas in the European honey
107 bees on this island have a low (~1E+03) viral load, and diverse range of DWV variants [2]. This
108 provides further evidence that DWV may be a general hymenopteran or insect virus rather than a
109 honey bee pathogen that has spilled over into the pollinator community. Again, the ubiquitous
110 presence of the DWV-C variant in *M. subnitida* colonies, and rarity in *A. mellifera* colonies on the
111 mainland again suggests limited viral exchange between these two species. The chance of spill-
112 over may be reduced due to the low (8E+07) DWV viral loads present in both the stingless and
113 honey bees of NE Brazil, relative to those found in asymptomatic (2.4E+09) and symptomatic
114 (6.9E+11) European honey bees [23]. Whereas, when these high DWV loads are present in honey
115 bees, DWV appears to spill-over into the neighbouring wasps and solitary bees [7]. These low
116 DWV viral loads in Brazil may be attributed to the hygienic habit of stingless bees [24], and
117 Varroa-tolerance in Africanised bees, both which will reduce the viral load in a colony.

118

119 **METHODS**

120 **Samples**

121 Pools of 30 *M. subnitida* workers were collected using a pooter directly at the entrance of
122 24 colonies from meliponiparies at ten mainland locations across NE Brazil. Samples from
123 Fortaleza and Mossoro were collected in 2016 with all other samples collected in 2013. In addition,
124 pools of ten *M. subnitida* workers from nine colonies located on the remote oceanic island of
125 Fernando de Noronha were collected in 2013 using the same method. These samples are interesting
126 since this population was originally established from 30 colonies brought to the island in 1983
127 from the mainland states of Ceara and Rio Grande do Norte [25]. In 1984 Kerr also established a
128 small population of European honey bees on Fernando de Noronha that were accidentally infested
129 by the Varroa mite, although the typically high levels of DWV were not present in either the honey
130 bees or Varroa [26].

131 During the same period pools of 30 healthy adult worker Africanised honey bees were
132 collected from the brood area of 12 colonies from six states across NE Brazil. All samples were
133 collected in absolute ethanol and stored at -20° C before transportation to the UK under license to
134 be analysed.

135

136 **Detection and quantification of DWV variants**

137 Total RNA was extracted from a pool of 10 heads per colony for both stingless and honey
138 bees. Heads were used as this reduced sample processing and is based on sound scientific

139 reasoning [27-30]. The heads were ground in liquid Nitrogen into a fine homogeneous powder, a
140 30mg sub-sample had its RNA extracted using a Qiagen RNeasy mini kit, which was enhanced by
141 using a QIAshredder kit for the *M. subnitida* samples [31]. Nanodrop (8000 series) quantification
142 was used to standardise the amounts of total RNA to 50 ng/μl using RNase free water, before been
143 stored at -80° C.

144 In order to quantify the viral load of each DWV Master-variant we used a recently
145 developed method [22]. Briefly, cDNA was synthesised using one-step SensiFAST SYBR No
146 ROX One-step kit (Bioline, London, UK), the reactions contained 1μl RNA at a concentration of
147 50 ng/μl, 10μl Senifast mix, 0.2μl Reverse transcriptase, 0.4μl RNase inhibitor, 0.75 pmol of each
148 primer (DWV F and R-Type A, B and C [Table 2]) and 7.5μl of H₂O. Reactions were run on a
149 Rotor-Gene Q Thermocycler (Qiagen) with an initial reverse transcription stage at 45° C for 10
150 min and a denaturation step of 95° C for 10 min, followed by 35 cycles of denaturation for 15 s at
151 95° C, annealing for 15 s at 58° C for primers A and B, and 61.5° C for primer C and extension
152 for 15 s at 72° C. A final dissociation melt curve was performed between 72° C and 90° C, at 0.5°
153 C increments, each with a 90 s hold. The melt curve was used to ensure that a single targeted
154 product was amplified, and that no contamination was present in the reverse transcription negative
155 controls or in the no-template controls. The threshold cycle (Ct) value was determined for each
156 sample using the QIAGEN Rotor—Gene Q Series Analysis software and viral quantification was
157 done by using serial dilutions of the standard DWV RNA, ranging from 1E+02 to 1E+07 copies
158 of DWV per reaction. All samples were run in triplicate and the average taken. Those samples
159 which had a standard deviation of ≥3 Ct were repeated. Furthermore, PCR products were run on a
160 2% agarose gel stained with 0.001% GelRed to confirm the correct sized band had been amplified.
161 A control housekeeping gene β-actin [23] was also run to ensure no degradation of the samples
162 had occurred, due to large distances these samples were transported both within and between
163 countries. Genome equivalents were calculated per sample using the following equation:

$$\text{Genome equivalents} = (\text{average copy number}) \times (\text{RNA dilution factor}) \times (\text{elution volume of RNA}) \times (\text{proportion of bee material})$$

166
167

168 **Funding information**

169 F. S. de Souza was funded by Empresa Brasileira de Pesquisa Agropecuária (EMBRAPA) (Cod.
170 SEG 02.11.01.029.00.00), CNPq (Proc. 400425/2014-9 and 305885/2017-0) and CAPES for the
171 PDSE scholarship (Proc. 88881.132120/2016-01).S. J. Martin and C. A. L. de Carvalho were
172 funded by a CNPq Special Visiting Researcher PVE (400425/2014-9).

173

174 **Acknowledgement**

175 The authors would like to acknowledge the Brazilian Institute of Environment and Renewable
176 Natural Resources, Chico Mendes Institute for Biodiversity Conservation, DAI/DEFN and
177 ADEFN, for the permission to conduct the survey in the island of Fernando de Noronha. We would
178 like to thank Márcia de Fátima Ribeiro, Candida B. da Silva Lima, Maria Emilene Correia de
179 Oliveira and Victor Hugo P. Dias for the collection of bee samples and the beekeepers, for allowing
180 the sampling of bees from their colonies. Also we would thank Bárbara Lins Caldas de Moraes for
181 support with the map.

182

183 **Author Contributions**

184 Conceptualization: All authors

185 Data curation: Flaviane S. de Souza, Jessica L. Kevill,

186 Formal analysis: Flaviane S. de Souza, Jessica L. Kevill, Stephen J. Martin.

187 Funding acquisition: Carlos A. L. de Carvalho, Stephen J. Martin.

188 Investigation: Flaviane S. de Souza, Jessica L. Kevill,

189 Methodology: Jessica L. Kevill,

190 Project administration: Carlos A. L. de Carvalho, Stephen J. Martin.

191 Resources: Carlos A. L. de Carvalho, Stephen J. Martin.

192 Supervision: Carlos A. L. de Carvalho, Stephen J. Martin.

193 Validation: Flaviane S. de Souza, Jessica L. Kevill,

194 Visualization: Flaviane S. de Souza

195 Writing - original draft: Flaviane S. de Souza

196 Writing - review & editing: Flaviane S. de Souza, Jessica L. Kevill, Stephen J. Martin.

197

198

199 **Conflicts of interest**

200 The authors declare that there are no conflicts of interest.

201

202 **Ethical statement**

203 There are no ethical issues.

204

205

206

207

208

209 **References**

- 210
- 211 **1. Rasmussen C, Cameron SA.** Global stingless bee phylogeny supports ancient divergence,
212 vicariance, and long-distance dispersal. *Biol J Linn Soc* 2010: 99:206–232
- 213 **2. Wille A.** Biology of the stingless bees. *Ann Rev Entomol* 1983: 28:41–64.
- 214 **3. Jaffé R, Pope N, Carvalho AT, Maia UM, Blochtein B et al.** Bees for Development:
215 Brazilian Survey Reveals How to Optimize Stingless Beekeeping. *PLoS ONE*. 2015: 10(3):
216 e0121157
- 217 **4. Nascimento VA, Matusita SH, Kerr WE.** Evidence of hybridization between two species of
218 *Melipona* bees. *Genetic Mol Biol* 2000: 23:79-81.
- 219 **5. Winston ML.** *Killer Bees: The Africanized Honeybee in The Americas*. 1992. Cambridge,
220 Mass.: Harvard University Press. ISBN 0-674-50353-8.
- 221 **6. Rosenkranz P, Aumeier P, Ziegelmann B.** Biology and control of *Varroa destructor*. *J*
222 *Invert Path* 2010: 103 Suppl 1:S96-119. doi: 10.1016/j.jip.2009.07.016
- 223 **7. Santamaria J, Villalobos EM, Brettell LE, Nikaido S, Graham JR, Martin SJ.** Evidence
224 of Varroa-mediated Deformed Wing virus spillover in Hawaii. *J Invert Path* 2018: 151:126-130
- 225 **8. Singh R, Levitt A, Rajotte E, Holmes E, Ostiguy N et al.** RNA viruses in Hymenopteran
226 pollinators: evidence of inter-taxa virus transmission via pollen and potential impact on non-*Apis*
227 Hymenopteran species. *PLoS ONE* 2010: 5(12):1-16.
- 228 **9. Evison SE, Roberts KE, Laurenson L, Pietravalle S, Hui J et al.** Pervasiveness of
229 Parasites in Pollinators. *PLoS ONE* 2012: 7(1) e30641.
- 230 **10. Levitt AL, Singh R, Cox-Foster DL, Rajotte E, Hoover K, Ostiguy N, Holmes EC.**
231 Cross-species transmission of honey bee viruses in associated arthropods. *Virus Res* 2013:
232 176(1):232-240.
- 233 **11. Manley R, Boots M, Wilfert L.** Emerging viral disease risk to pollinating insects:
234 ecological, evolutionary and anthropogenic factors. *J Appl Ecol* 2015: 52(2):331-340.
- 235 **12. Mazzei M, Carrozza ML, Luisi E, Forzan M, Giusti M et al.** Infectivity of DWV
236 Associated to Flower Pollen: Experimental Evidence of a Horizontal Transmission Route. *PLoS*
237 *ONE* 2014: 9(11): e113448.
- 238 **13. Tehel A, Brown MJ, Paxton RJ.** Impact of managed honey bee viruses on wild bees. *Curr*
239 *Opinion Virol* 2016: 19:16-22.

- 240 **14. Li J, Peng W, Wu J, Strange JP, Boncristiani H, Chen Y.** Cross-species infection of
241 Deformed wing virus poses a new threat to pollinator conservation. *J Eco Entomol* 2011:
242 104(3):732-739.
- 243 **15. Fürst, M, McMahon D, Osborne J, Paxton R, Brown, M.** Disease associations between
244 honeybees and bumblebees as a threat to wild pollinator. *Nature* 2014: 506: 364-366.
- 245 **16. Graystock P, Goulson, D, Hughes WO.** Parasites in bloom: flowers aid dispersal and
246 transmission of pollinator parasites within and between bee species. *Proc Roy Soc B Biol Sci*
247 2015: 282:20151371
- 248 **17. Alvarez LJ, Reynaldi FJ, Ramello PJ, Garcia MLG, Sguazza GH et al.** Detection of
249 honey bee viruses in Argentinian stingless bees (Hymenoptera: Apidae) *Insect. Soc* 2017:
250 65:191-197.
- 251 **18. Ueira-Vieira C, Almeida LO, de Almeida FC, Amaral IMR, Brandeburgo AM, Bonetti**
252 **AM.** Scientific note on the first molecular detection of the acute bee paralysis virus in Brazilian
253 stingless bees *Apidologie* 2015: 46: 628.
- 254 **19. Wilfert L, Long G, Leggett HC, Schmid-Hempel P, Butlin R et al.** Deformed wing virus
255 is a recent global epidemic in honeybees driven by Varroa mites. *Science* 2016: 351: 594–597.
- 256 **20. Ryabov EV, Childers AK, Chen Y, Madella S, Nessa A, van Engelsdorp JD.** Recent
257 spread of Varroa destructor virus-1, a honey bee pathogen, in the United States. *Sci Rep* 2017: 7:
258 17447.
- 259 **21. McMahon DP, Natsopoulou ME, Doublet V, Furst M, Weging S et al.** Elevated virulence
260 of an emerging viral genotype as a driver of honeybee loss. *Proc Roy Soc B Biol Sci* 2016:
261 283:20160811.
- 262 **22. Kevill JL, Highfield A, Mordecai GJ, Martin SJ, Schroeder DC.** ABC Assay: Method
263 Development and Application to Quantify the Role of Three DWV Master Variants in
264 Overwinter Colony Losses of European Honey Bees. *Viruses* 2017: 9: 314.
- 265 **23. Highfield AC, Nagar AE, Mackinder LC, Noel LM, Hall MJ et al.** Deformed wing virus
266 implicated in overwintering honeybee colony losses. *Appl Environ Microbiol* 2009: 75: 7212-
267 7220.
- 268 **24. de Jesus JN, Chambó ED, da Silva-Sodré G, Oliveira NTE, Carvalho CA.** Hygienic
269 behavior in *Melipona quadrifasciata anthidioides* (Apidae, Meliponini). *Apidologie* 2017:
270 48:504-512.

- 271 **25. Kerr WE, Cabeda M.** Introdução de abelhas no território federal de Fernando de Noronha.
 272 *Rev of Ciencia e Cultura* 1985: 37: 467-471.
- 273 **26. Brettell LE, Martin SJ.** Oldest Varroa tolerant honey bee population provides insight into
 274 the origins of the global decline of honey bees. *Sci Rep* 2017: 7:45953.
- 275 **27. Fujiyuki, T, Takeuchi H, Ono M, Ohka S, Sasaki TS et al.** Novel Insect Picorna-Like
 276 Virus Identified in the Brains of Aggressive Worker Honeybees. *J. Virol* 2004: 78: 1093-1100.
- 277 **28. Yue C. Genersch E.** RT-PCR analysis of Deformed wing virus in honeybees (*Apis*
 278 *mellifera*) and mites (*Varroa destructor*). *J Gen Virol* 2005: 86(12): 3419-3424.
- 279 **29. Genersch E, Von Der Ohe W, Kaatz H, Schroeder A, Otten C et al.** The German bee
 280 monitoring project: a long term study to understand periodically high winter losses of honey bee
 281 colonies. *Apidologie* 2010: 41(3):332-352.
- 282 **30. Zioni N, Soroker V, Chejanovsky N.** Replication of Varroa destructor virus 1 (VDV-1)
 283 and a Varroa destructor virus 1–deformed wing virus recombinant (VDV-1–DWV) in the head
 284 of the honey bee. *Virology* 2011: 417(1):106-112.
- 285 **31. Forsgren E, Locke B, Semberg E, Laugen AT, Miranda JR.** Sample preservation,
 286 transport and processing strategies for honeybee RNA extraction: Influence on RNA yield,
 287 quality, target quantification and data normalization. *J Virol Meth.* 2017: 246:81-89.

288
 289
 290
 291
 292
 293 **Table 1.** The mean viral load of each DWV master variant detected in the 21 *Melipona subnitida*
 294 and 12 *Apis mellifera* samples collected from across NE Brazil.

	<i>Melipona subnitida</i>	<i>Apis mellifera</i>
	Average viral load	Average viral load
DWV-A	8.10E+07	6.96E+07
DWV-B	n.d.	2.35E+05
DWV-C	7.31E+06	2.06E+06
All	8.83E+07	7.19E+07


296
 297 **Table 2.** Primers used in this study were developed by [24].

Target	Primer Name	Sequence (5' - 3')	Size of product (bp)
DWV Forward	DWVnew-F1	TACTAGTGCTGGTTTTTCCTTT	299
DWV Type A	DWVA-R1	CTCATTAACTGTGTCGTTGAT	155
DWV Type B	DWVB-R1	CTCATTAACTGAGTTGTTGTC	155
DWV Type C	DWVC-R1	ATAAGTTGCGTGGTTGAC	152

303

304


305



306

307

308 **Fig. 1.** Typical gel showing the presence of β -actin in all samples of *Melipona subnitida*, *Apis*
 309 *mellifera* and positive controls, confirming that the samples contained intact RNA.



310
 311
 312 **Fig. 2.** Proportion and viral load of DWV-A (red), B (blue) and C (green) variants detected in A)
 313 *Melipona subnitida* stingless bees and B) *Apis mellifera* from across NE Brazil. The sample
 314 locations are 1. Cumaru, 2. Exu, 3. Fortaleza, 4. Fernando de Noronha, 5. Mata Grande, 6.
 315 Mossoró, 7. Passira, 8. Paulo Afonso, 9. Riacho das Almas, 10. Taquaritinga do Norte, 11. Água
 316 Branca, 12. Piranhas, 13. Cruz das Almas, 14. Seabra, 15. Garanhuns 16. Mossoró, 17. São
 317 Cristóvão, 18. Areal. The states are CE= Ceara, RN= Rio Grande Do Norte, PB= Paraíba, PE=
 318 Pernambuco, AL= Alagoas, SE= Sergipe and BA= Bahia.
 319