



## 6 **Abstract**

7

8 **Objectives:** Isometric muscle contractions are used in the management of patellar tendinopathy to  
9 manage symptoms and improve function. Little is known about whether long or short duration  
10 contractions are optimal and the mechanisms. This study examined the immediate and short term  
11 effects of long and short duration (four weeks) isometric contraction on patellar tendon pain,  
12 quadriceps muscle function and tendon adaptation.

13 **Design:** Repeated measures within subjects

14 **Methods:** Sixteen participants with patellar tendinopathy were randomised to either short duration (24  
15 sets of 10 seconds) or long duration (six sets of 40 seconds) isometric loading, performed on a leg  
16 extension machine, five times per week for four weeks. Loading was performed at estimated 85%  
17 maximal voluntary contraction (MVC). Pain on single leg decline squat and tendon adaptation (change  
18 in tendon thickness both over the weeks and within session) were assessed at baseline, two and four  
19 weeks. Prior to the 4-week study, a sub-sample of 8 participants performed both protocols in a random  
20 order 5-7 days apart to determine immediate pain response on hop and single leg decline squat tests.

21 **Results:** There was a significant reduction in pain following isometric loading on both SLDS ( $p < 0.01$ )  
22 and hop tests ( $p < 0.01$ ). Pain improved from weeks two to four ( $p < 0.05$ ), and quadriceps strength  
23 improved from baseline to week two and week two to week four ( $p < 0.05$ ). There were no group or  
24 interaction effect for immediate change in pain, or change in pain and quadriceps strength over four  
25 weeks. Tendon thickness did not change over the study period. There was significant transverse strain  
26 at each measurement occasion ( $P < 0.01$ ), although there were no group or interaction effects.  
27 Percentage transverse change did not change across the three measurement occasions ( $p = 0.08$ ).

28 **Conclusions:** This is the first study to show that short duration isometric contractions are as effective  
29 as longer duration contractions for relieving patellar tendon pain when total time under tension is  
30 equalised. Improvements in pain over the first four weeks of isometric loading parallel improved  
31 strength, but there was no evidence of tendon adaptation over the short 4 week study period. This  
32 finding provides clinicians with greater options in prescription of isometric loading and may be  
33 particularly useful among patients who do not tolerate longer duration contractions.

34

35 Key words: patellar tendinopathy, isometric loading, rehabilitation, ultrasound, biomechanics,

36 transverse strain

37

38

## 39 **Introduction**

40 Patellar tendinopathy is characterised by localised pain and pathology at the inferior pole of the patella  
41 that is common among jumping athletes and often impairs performance. Isometric loading has recently  
42 become more popular in managing patellar tendon pain (Rio et al., 2015, Cook and Purdam, 2014).  
43 Longer duration isometric contractions (45 seconds) of high intensity (70% maximal voluntary  
44 contraction (MVC)) are commonly used as preliminary evidence suggests they offer a greater  
45 immediate reduction in pain than isotonic loading in patellar tendinopathy (Rio et al., 2015). Isometric  
46 exercises are therefore recommended in managing patellar tendinopathy pain among competing or ‘in-  
47 season’ athletes. Isometric loading is also recommended at the commencement of rehabilitation for  
48 patellar tendinopathy if isotonic exercises are too painful (Malliaras 2015).

49

50 To date, no studies have compared the clinical effects of short and long duration isometric loading.  
51 Relatively long duration contractions of 45 seconds are recommended and have been shown to have  
52 immediate effects on pain in patellar tendinopathy (Rio et al. 2015). Isometric loading has been  
53 shown to lead to immediate reversal of motor cortex inhibition and increased motor output (MVC) and  
54 this may be secondary to immediate pain reduction following this exercise intervention in patellar  
55 tendinopathy (Rio et al., 2015).

56

57 Shorter duration contractions have been shown to have an effect on tendon adaptation (Arampatzis  
58 2010) but their efficacy in regards to managing pain in patellar tendinopathy is not known. Therefore,  
59 investigating the relative efficacy of short and long duration contractions in managing patellar  
60 tendinopathy is warranted.. It is generally agreed that there is altered tendon morphology in  
61 tendinopathy, characterised by an increase in glycosaminoglycan (GAG) content in tendinopathic  
62 tendons (Maffulli et al., 2006). These characteristic changes in tendon structure can result in  
63 reductions in fluid exchange via reduced tissue fluid permeability, and possibly reduced free water  
64 content (Henninger et al., 2010; Dubinskaya et al., 2007). In support of this, reduced tendon thickness  
65 immediately following a loading protocol (‘transverse strain’ related to fluid exchange) has been  
66 shown to occur in healthy tendons following loading, but the response is blunted in tendinopathic

67 tendons (Grigg et al., 2012; Wearing et al., 2013). No studies have investigated how the transverse  
68 strain response (change in thickness) changes over the course of loading rehabilitation for patellar  
69 tendinopathy. This tendon adaptation outcome may help to improve understanding of loading  
70 associated mechanisms in tendinopathy and is favoured to subjective measures of tendon structure on  
71 ultrasound and MRI that demonstrate poor repeatability.

72

73 No previous study has compared the effects of long and short duration isometric loading on clinical  
74 (pain) and tendon adaptation (transverse strain, tendon thickness) outcomes during the initial phase of  
75 rehabilitation for patellar tendinopathy. Therefore the aims of this study are; (i) to investigate the  
76 immediate effects of long and short duration isometric loading on pain and tendon adaptation (tendon  
77 thickness, transverse strain) in patellar tendinopathy; (ii) to investigate the effects of long and short  
78 duration isometric loading on pain and tendon adaptation outcome in the initial phase (time course of  
79 4 weeks) of rehabilitation for patellar tendinopathy.

80

## 81 **Methods**

### 82 **Participants**

83 Sixteen males with unilateral or bilateral patellar tendinopathy participated in the study. Participants  
84 were recruited from State level Volleyball and Basketball leagues, as well as through local advertising  
85 and word-of-mouth. Participants were included if they had pain localized to the inferior patella pole  
86 that was aggravated by jumping, and had ultrasound imaging confirmed patellar tendon pathology  
87 (hypoechoic regions on gray scale +/- Doppler signal). Participants had to be willing to attend a gym  
88 to perform isometric loading exercise five times per week for four weeks, and stop jumping and  
89 running activities during the four-week study period. Potential participants were excluded if they had  
90 had previous patellar tendon surgery or rupture, other diagnoses that could explain their pain (e.g.  
91 Hoffa's fat pad syndrome, patellofemoral joint pain), rehabilitation or injections for their patellar  
92 tendon pain in the previous three months, or any other lower limb injury that would prevent them  
93 completing the rehabilitation loading. The study was approved by the local Human Ethics Committee.

94 Participants signed an informed consent form prior to inclusion into the study. The study conformed to  
95 the principles of the World Medical Associations Declaration of Helsinki.

96

### 97 **Pre-testing**

98 All testing was performed at a private physiotherapy clinic in Melbourne, Australia. At the initial visit  
99 participants completed a questionnaire that included demographic data (age, height, weight, sport  
100 played, leg dominance, whether they performed lower body weight training) and basic clinical  
101 information (side of patellar tendon pain, duration of pain - see table 1). Pain and function were  
102 assessed with the Victorian Institute of Sport Assessment for Patellar Tendinopathy (VISA-P), a  
103 reliable and valid questionnaire for patellar tendinopathy (Visentini et al., 1998). A score of 0  
104 represents the worst possible symptoms and function and a score of 100 represents no symptoms and  
105 full function. The VISA was performed so that baseline self-reported pain/dysfunction could be  
106 compared between the groups. This measure was not assessed over time because a large component  
107 (40%) relates to sporting function and in this study participants were asked to stop playing sport.

108

### 109 **Isometric loading interventions**

110 Participants were randomly allocated in blocks of six (selected a group number from an opaque  
111 envelope) to one of two groups: (i) short duration isometrics; or (ii) long duration isometrics. The  
112 short duration group performed 24 sets of 10 second isometric contractions with a 20 second rest  
113 between each repetition. The long duration group performed six sets of 40 second isometric  
114 contractions with an 80 second rest between each contraction. The total time under tension (240  
115 seconds or four minutes) and work to rest ratio (1:2) were the same for both groups. Isometric loading  
116 was performed on a leg extension machine at a knee flexion angle of 30 degrees (knee extension = 0°).  
117 This is usually a tolerable position to hold heavy load for someone with patellar tendinopathy.  
118 Participants were provided with custom made plywood cut at 30 degrees and they used this to ensure  
119 the appropriate knee angle during the home loading session.

120

121 Participants were required to perform the loading program five times per week over the four-week  
122 study period, this was performed at the participants' gym (if they had a gym membership) or the  
123 physiotherapy clinic. During the rest time for one side participants were instructed to load the opposite  
124 side using an identical protocol, so they performed the leg extension loading unilaterally, but on both  
125 sides. Participants were asked to abstain from any form of weight-bearing exercise that loads the knee  
126 extensors, including running, hopping, jumping, squatting, or any lower body weights/strength  
127 exercises. They were also asked not to perform an isometric exercise session within 24 hours of their  
128 follow up appointments.

129

### 130 **85% MVC testing**

131 At baseline, week 2 and week 4, isometric loading was performed in order to estimate 85% MVC since  
132 tendon adaptation appears to occur with loads >70% MVC (Bohm et al. 2015). This was estimated  
133 during on a leg extension machine (Impulse Fitness, Newbridge, Scotland) during an isometric hold at  
134 an angle of 30 degrees knee flexion. A laser pointer was attached to the moving arm of the leg  
135 extension machine so that when the knee was in 30 degrees (measured with a goniometer) the laser  
136 pointer was aimed at a mark that was placed on a whiteboard in front of the machine. The affected  
137 side or worse side (among bilateral participants) was tested. None of the participants reported pain that  
138 was more than minimal (defined as 3/10) during the leg extension loading sessions. Participants were  
139 required to perform an isometric leg extension hold for 40 seconds at this angle. Immediately after the  
140 task, perceived exertion was rated on a Borg scale from 0 to 20 (Borg 1982). A rating of 17-18 was  
141 estimated to correspond to 85% MVC. If the rating was lower, the task was repeated with 5kg  
142 additional load. If the rating was higher or 40 seconds was not reached, the task was repeated with 5kg  
143 less load. This was repeated until 85% MVC for a 40 second isometric contraction was estimated,  
144 generally within two to three trials. This represents a pragmatic, clinic-based method of assessing  
145 appropriate load intensity. Participants rested for two minutes between trials. Participants were  
146 instructed to monitor their rating of perceived exertion and increase the load if their perceived exertion  
147 was lower than 17/20 on the Borg scale during any home loading session.

148

149 **Tendon thickness and transverse strain testing**

150 Transverse strain was defined as the percentage reduction in patellar tendon thickness following an  
151 isometric loading protocol. Pilot testing revealed that a very short duration isometric protocol (10  
152 repetitions of four second contraction with a four second rest between contractions, and repeating six  
153 sets with a one-minute rest between sets) produced greater immediate transverse strain than the  
154 loading interventions described above (i.e. repeated 10 or 40 second contractions), and was less  
155 fatiguing, less likely to interfere with the training protocols and more clinically practical. Therefore,  
156 we used this isometric protocol to assess (i) immediate post loading transverse strain; and (ii) short  
157 term (comparing baseline to four weeks) transverse strain for each loading intervention. The training  
158 load used during transverse strain assessment was the 85% MVC estimated load described above.  
159 Isometric loading was performed on a leg extension machine in 30 degrees of knee flexion. As  
160 described for the home loading protocols, a laser point was used to ensure the knee angle was  
161 maintained.

162  
163 Ultrasound imaging measurements were performed at baseline and after each of the six sets which  
164 consisted of 10 repetitions of 4 second isometric contractions, producing seven measurement  
165 occasions in total. Participants were lying supine on a treatment plinth with the knee that was being  
166 imaged flexed at 90 degrees. Patellar tendon thickness was measured using an ultrasound machine  
167 with a 12 MHz linear array transducer (Mindray M7, Mindray, Shenzhen, China) set at a depth of  
168 three cm. Minimal pressure was applied to the skin to avoid compressing the tendon with the probe.  
169 The proximal thickness of the tendon (10mm distal to the inferior patellar pole) was measured with the  
170 ultrasound probe placed in the sagittal plane. The centre point of the patellar tendon insertion into the  
171 inferior pole of the patella was measured with ultrasound and marked on the skin with a pen. Sagittal  
172 plane images were recorded, with the centre of the probe aligned with the pen mark. Care was taken to  
173 ensure some of the patella (bone) appeared in the recorded image (Figure 1) and the probe was aligned  
174 perpendicular to the tendon. Three ultrasound images were recorded at each measurement occasion  
175 and the mean thickness was used in analysis. Overall, there were two outcomes used in analyses: 1)  
176 Resting tendon thickness (this was taken as the baseline measure for that particular session prior to



177 any loading); 2) transverse strain (percentage change in thickness from resting to post loading  
178 protocol).

179

180 Figure 1. Typical sagittal image showing the inferior patellar pole and patellar tendon.

181

182 Tendon thickness assessment was performed by one of the researchers (SS) who was trained by an  
183 experienced ultrasonographer and had over 30 hours of practice prior to commencing testing. The  
184 ultrasonographer was not blind to group allocation but measurement of tendon thickness from the still  
185 images was performed by a blinded assessor. Intra-rater reliability of tendon thickness measures were  
186 assessed among a subset of eight participants. Tendon thickness was measured on four occasions for  
187 each participant, prior to which no loading of the tendon was carried out. There was a two-minute  
188 break between testing during which participants stood up and were then repositioned again on the  
189 treatment plinth. Intra-rater reliability was estimated using interclass correlation coefficients (ICC (1,  
190 2) = 0.95, 95% CI = 0.87-0.99). The minimal detectable change (95% confidence) was 0.17 mm.

191

### 192 **Follow up assessment**

193 Participants were followed up at two and four weeks. On both occasions, 85% MVC during the 40  
194 second isometric hold was reassessed and tendon thickness, transverse strain were measured using  
195 identical procedures already outlined. Participants were also asked how many of the five weekly  
196 sessions they completed in order for exercise adherence to be calculated (percentage of sessions  
197 completed = (completed sessions/scheduled sessions)x100). Abstinence from activity involving  
198 impact loading for the knee, such as running and jumping, was also assessed. Pain during a single leg  
199 squat was used to monitor changes in pain during the four-week intervention. Participants performed a  
200 standard test used to elicit patellar tendon pain, involving a single leg squat to 60 degrees knee flexion  
201 whilst standing on a decline board (Zwerver et al., 2007). Pain intensity during the test was rated using  
202 a 100 mm visual analogue scale.

203

### 204 **Immediate pain response following isometric loading**

205 Prior to commencement of the study, a subsample of eight participants were included in a randomised  
206 cross-over study investigating the immediate effects of the isometric protocols on pain in patellar  
207 tendinopathy. Participants were randomised to perform either the short duration (24 x 10 second  
208 isometric contractions with a 20 second rest between contractions) or long duration (6 x 40 second  
209 isometric contractions with an 80 second rest between contractions) protocol. They performed the  
210 second protocol on a separate occasion within 5-7 days. Five repetitions of two functional tests were  
211 performed before and after the isometric loading. These included a single leg squat to 60 degrees knee  
212 flexion (Zwerver et al., 2007) and a single leg submaximal hop (participants were instructed to hop  
213 continuously with hands on hips). Pain intensity during the functional tests was rated using a 100 mm  
214 visual analogue scale.

215

#### 216 **Data processing and reliability**

217 All images were exported into jpeg format for determination of tendon thickness (Image j - Wayne  
218 Rasband National Institute of Health, Bethesda, MD, USA). The images were calibrated to enable a  
219 pixel to mm ratio to be determined. Measures of patellar tendon thickness (anterior to posterior) were  
220 then made at a distance of 10mm distal to the inferior patellar pole (this site was chosen as patellar  
221 tendon pathology typically occurs at the proximal tendon).

222

#### 223 **Data analysis**

224 Statistical analysis was performed using SPSS (version 22, SPSS Inc., Chicago, IL). Baseline group  
225 characteristics were compared using independent t-tests (age, height, weight, duration of symptoms,  
226 VISA, patellar tendon AP thickness, leg extension 85% MVC). Change in pain during the single leg  
227 decline squat task, leg extension 85% MVC and patellar tendon thickness were assessed over the four  
228 weeks, and compared between the two groups (Two way (Gp x wks) mixed model analysis of variance  
229 ANOVA). Change in transverse strain (acute change in tendon thickness from rest) was assessed  
230 within each test session and between each group (Two way (Gp x sess) mixed model ANOVA).  
231 Transverse strain at each test occasion was also compared between the three test occasions (repeated  
232 measures ANOVA). For the cross-over repeated measures study, immediate change in pain during the

233 single leg decline squat and hop tasks were assessed following each loading protocol and compared  
234 between the groups (Two way (Gp x sess) mixed model ANOVA).

235

## 236 **Results**

237 Sixteen men with patellar tendinopathy were randomized into the groups (short duration (n=8); long  
238 duration (n=8)). There were no significant baseline differences between the groups for demographic  
239 characteristics (age, height, weight), duration of symptoms, VISA-P score, patella tendon AP  
240 thickness or 85% MVC (Independent t-tests,  $p>0.05$ ) (Table 1). The mean VISA-P scores (56 to 62)  
241 indicated that participants' sporting activity was limited by their patellar tendon pain. Other  
242 characteristics including weight training performed, leg dominance, and bilateral pain were also  
243 similar between the groups (Table 1).

244

245 All participants were active in sports involving impact loading of the knee (Table 2). All participants  
246 had previously undertaken load-based rehabilitation for their patellar tendon pain for a minimum of  
247 three months. In both groups, seven (87.5%) participants adhered to no sport participation during the  
248 four-week study period. Almost all participants reported that they completed all of the prescribed five  
249 sessions per week (96% in the low duration and 100% in the long duration group).

250

251 Table 1: Characteristics of the two groups at baseline

	<b>short duration</b>	<b>long duration</b>
	<b>(n=8)</b>	<b>(n=8)</b>
Age (yrs)	26 (4.4)	30 (4.1)
Height (cm)	184.5 (5.6)	186.8 (7.3)
Weight (kg)	84.5 (9.5)	87.1 (9.4)
Duration of symptoms (yrs)	2.3 (2.0)	3.4 (1.9)
VISA-P	53 (14.5)	58 (12.8)
AP thickness (mm)	5.6 (1.7)	6.1 (1.0)

85% MVC (kg)	38.4 (6.3)	41.0 (5.6)
Bilateral pain	1 (12.5)	2 (25.0)
Right side injured/worst	6 (75)	5 (62.5)
Right leg dominance	7 (87.5)	7 (87.5)
Gym weights	6 (75.0)	6 (75.0)
Manual work	0 (0.0)	1 (12.5)

252 NB: median (interquartile range) for parametric, and frequency (proportion) for discrete data

253

254

255 Table 2: Sport played by the participants

	short duration	long duration
Football	2 (25)	5 (63.5)
Basketball	2 (25)	1 (13.5)
Volleyball	1 (13.5)	none
Cricket	1 (13.5)	1 (13.5)
Running	1 (13.5)	none
Cheerleading	1 (13.5)	none
Rock climbing	1 (13.5)	none
Rugby	none	1 (13.5)

256

257 **Immediate change in pain**

258 There was a significant reduction in pain following isometric loading on both SLDS ( $p < 0.01$ ) and hop  
259 tests ( $p < 0.01$ ) (Figure 2). There was no significant difference between long or short duration isometric  
260 loading for either SLDS ( $p = 0.95$ ) or hop ( $p = 0.75$ ) and no significant interaction effects (SLDS,  
261  $p = 0.60$ ; hop,  $p = 0.33$ ).

262 For the SLDS test, one participant experienced an increase in pain, two had no change and the  
263 remaining reported reduced pain. The mean change on the SLDS test was small (mean = 0.8, SD = 0.9,

264 range = -0.2 to 3.0). All participants experienced reduced pain on the hop test and again the mean  
265 change was small (mean = 1.7, SD = 3, range = 0.1 to 4.3). There was no difference in pain changes  
266 scores among participants grouped as having high versus low baseline pain scores (SLDS,  $p=0.60$ , hop,  
267  $p = 0.33$ ).

268

269 Figure 2. Immediate effects of long and short duration isometric loading on pain on single leg decline  
270 squat and hop tests

271

### 272 **Change in pain over 4 weeks**

273 Pain reported during single-leg decline squat testing reduced significantly during the study period but  
274 only between week two and four (Figure 3). There was no significant difference between long or short  
275 duration isometric loading ( $p=0.34$ , power = 0.15) or interaction effect for change in pain on the SLDS  
276 test ( $p=0.65$ , power = 0.11).

277

278 Figure 3. Pooled single leg decline squat (SLDS) pain scores across the training period. (Mean  $\pm$  SD).

279 \* Significantly differences between weeks ( $p < 0.05$ ).

280

### 281 **Change in strength and tendon outcomes over 4 weeks**

282 Eighty-five percent MVC increased significantly from baseline to two weeks, and from two to four  
283 weeks (Figure 4). There was no significant difference between long or short duration isometric loading  
284 ( $p=0.21$ , power = 0.23) or interaction effect for change in 85% MVC ( $p=0.11$ , power = 0.43).

285

286 Figure 4. Pooled values for 85% isometric loading at baseline, week 2 and week4 (Mean  $\pm$  SD). \*

287 Significantly differences between weeks ( $p < 0.05$ ).

288

289 Resting anteroposterior thickness did not change over the three sessions ( $p=0.26$ , power = 0.23) and  
290 there were no group (long or short duration loading) ( $p>0.66$ , power = 0.10) or interaction effects  
291 ( $p=0.58$ , power = 0.66). There was a significant reduction in AP thickness (i.e. transverse strain)

292 within each session immediately following the loading protocol ( $p < 0.01$ ) (see Figure 5) but there were  
293 no group ( $p > 0.90$ ) or interaction effects ( $p = 0.81$ ). Percentage tendon transverse strain was 14% or  
294 greater at each session (session 1 = 14%, session 2 = 17%, session 3 = 22%). Although transverse  
295 strain increased across the training period from ~14 % to 22%, this was not significant ( $p = 0.08$ ).

296 **Discussion**

297 Isometric loading is a popular modality for treating patellar tendon pain and long duration isometric  
298 contractions are recommended (Rio et al., 2015, Cook and Purdam, 2014). Our data suggests that long  
299 and short duration isometrics (with total time under tension equalised) have a similar effect on  
300 immediate pain among people with patellar tendinopathy. Further, we found that isometric contraction  
301 was associated with reduced pain, increased quadriceps strength but did not have a significant effect  
302 on patellar tendon thickness or transverse strain in the initial period of rehabilitation for patellar  
303 tendinopathy. There were no differences in pain and strength outcomes for the long and short duration  
304 isometric contractions. Our findings provide clinicians with greater options when approaching  
305 rehabilitation of patients with patellar tendinopathy. In some instances, isometric loading may be  
306 better tolerated with a shorter 10 second contraction time (Malliaras et al. 2015). Further work is  
307 needed to determine if reducing the total time under tension, which is likely to positively influence  
308 adherence, would lead to similar pain improvement.

309  
310 Recommendations of long duration isometric contractions is based on a single study among a  
311 comparable population (VISA-P mean 53 versus approximately 58 in our study) of active athletes with  
312 patellar tendinopathy (Rio et al. 2015). Rio et al. (2015) reported a pain reduction from a mean of  
313 seven to 0.17 (numeric rating scale from zero to ten) on single leg decline squat testing following 5  
314 repetitions of 45 seconds of isometric loading at 70% MVC. It can be argued that our load intensity of  
315 85% MVC of a 40 second hold is comparable to the Rio et al. short duration 70% (Brzycky 1993).  
316 Despite similar protocols, we found a more modest reduction in pain (3.4 to 2.6 on a visual analogue  
317 scale) during single leg decline squat testing. There are key study differences that may explain the  
318 discrepancy. First, Rio et al. performed loading at 60 degrees versus our protocol performed at 30  
319 degrees of knee flexion, so the load may have been higher in their study. Second, even though VISA-P  
320 scores were comparable between the groups, the Rio et al. (2015) cohort had greater pain on single leg  
321 decline squat testing at baseline (mean of 7 versus mean of 3 in the current study). Third, using the  
322 numerical rating scale may have biased participants towards larger and more definite changes in pain  
323 following the intervention.

324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351

As expected, given the loading program undertaken and cessation of aggravating activities such as jumping, pain with loading (single leg decline squat test) improved during the study between weeks two and four. As with immediate changes in pain, participants in both the long and short duration isometric groups experienced similar improvement in pain during the final two weeks of the intervention. Given we did not include a control group that rested from sporting activity but did not perform isometric loading, we do not know how much pain improvement relates to stopping sporting activity. It is likely that reducing sports activity had some effect. However, our pragmatic study was designed to replicate the initial few weeks of rehabilitation, when people are often advised to moderate sporting activity, and to specifically compare the two isometric protocols.

Participants in this study experienced significant improvement in leg extension strength following the heavy isometric loading, as would be expected following loaded exercise interventions, and there were no group differences in strength adaptation. Lack of group differences in this outcome is consistent with reports that the time under tension integral is the most important determinant of muscle size and strength adaptation (Riog 2009). In our study time under tension and load intensity (tension) was equalised between the groups.

In contrast to muscle strength adaptation, there was no change in tendon thickness or immediate transverse strain response to load across the four-week intervention. This supports the view that tendon adaptation most likely will take several months and lags behind neuromuscular adaptation (Bohm et al. 2015). Tendon is generally less responsive than muscle in the short term but it is important to note that we only considered two tendon adaptation outcomes and did not consider microstructural change (eg change in fibril morphology) that have been shown to change in patellar tendinopathy following an exercise intervention (Kongsgaard 2010). So although our data suggest no tendon adaptation following our isometric loading protocols, and no differences between groups, more data on other outcomes and longer follow-up times are needed.



352 The trend for increased transverse strain across the four weeks is worth mentioning. Here we show  
353 relatively large immediate reductions in tendon thickness (14%) with cyclic isometric loading, and  
354 although not significant, a large effect size indicating greater changes in tendon thickness reductions  
355 chronically across the sessions (22%). This may be explained by the increased load (85% MVC) under  
356 which the transverse strain test was performed at week two and four. However, we chose this design  
357 as it represents what occurs in clinical practice when exercise loading is increased progressively. Even  
358 if the trend towards an increased fluid flow response is related to increasing loading, it does suggest  
359 that the tendon matrix is responsive in the short term to heavier loading. Regardless, we did not see a  
360 decrease in thickness of the tendon over the four weeks suggesting that a transient tendon response to  
361 load does not translate to tendon adaptation, but this may be explained by our short follow up time.

362

363 Previous work reported that tendinopathy resulted in a blunted response of the tendon transverse strain  
364 with loading compared to healthy subjects. Using a combination of concentric and eccentric loading  
365 (double legged squats -45 reps at 145% body weight) Wearing et al. (2015) reported that transverse  
366 strain changes in tendinopathic tendons were minimal (0.2%), and significantly less than that of  
367 healthy tendons (~ 6%). In contrast to these findings, we showed that cyclic isometric loading  
368 produces a much greater transverse strain response. Wearing et al (2013) reported transverse strain  
369 values for healthy subjects in the patellar tendon of ~22.5 % with 90 repetitions of double legged  
370 squats, at a loading of 175% body weight. Thus, our finding of increased transverse strain over the  
371 four-week period being similar to that in healthy subjects is indeed interesting. However, the fact that  
372 we also show relatively large transverse strains in the initial stages suggests that there are differential  
373 stimuli using our isometric protocol to the previous dynamic protocols described. It could be that the  
374 'time under tension' is the major influencing factor. Tendon is described as being viscoelastic and as  
375 such exhibits strain in a time dependant manner. With this in mind, it can be understood that if a stress  
376 is applied for a longer period of time, then the tendon structure will likely affect the viscous element to  
377 a larger degree and undergo more strain (creep), all things being equal. This resultant increased  
378 longitudinal strain may also be accompanied by a concomitant reduction in tendon thickness. More

379 work is needed to understand the mechanisms underpinning change in patellar tendon transverse strain  
380 (fluid loss or other mechanism), and the relevance to clinical outcome in patellar tendinopathy.

381

382 This is the first study to show that there are beneficial effects on patellar tendon pain from both long  
383 and short duration isometric contractions, but there are study limitations that need to be highlighted. A  
384 control group of people with patellar tendinopathy that stopped sporting activity but did not perform  
385 isometric loading would have allowed us to delineate the effect of exercise versus rest from other  
386 activities on improved pain during the study. The sample size may have also limited power in  
387 identifying significant differences (e.g. change in transverse strain response from baseline to four  
388 weeks). In addition, a longer training period and increased participant numbers would add to the  
389 ability of this study to discriminate potential mechanistic changes exhibited by tendinopathic tendons  
390 with chronic isometric loading.

391

392 Isometric exercises are a popular treatment for patellar tendon pain, and long duration contractions are  
393 recommended. This study found that long and short duration isometrics contractions are equally  
394 effective for immediate relief of patellar tendon pain when total time under tension is equalised.  
395 Improvements in pain over the first four weeks of isometric exercise parallel improved strength, but  
396 there was no evidence of tendon adaptation over the short 4 week study period. This finding provides  
397 clinicians with greater options when prescribing isometric exercise for patellar tendinopathy.

398

399 Acknowledgments

400 The authors would like to thanks all participants involved in this study. There are no external sources  
401 of funding associated with this work.

402

403 Conflict of interest

404 None of the authors have any conflict of interest to declare in relation to this manuscript

405

406 References

407

408 Arya, S., Kulig, K., 2010. Tendinopathy alters mechanical and material properties of the Achilles  
409 tendon. *J. Appl. Physiol.* 108, 670–5. doi:10.1152/jappphysiol.00259.2009

410 Beyer, R., Kongsgaard, M., Hougs Kjær, B., Øhlenschläger, T., Kjær, M., Magnusson, S.P., 2015.

411 Heavy Slow Resistance Versus Eccentric Training as Treatment for Achilles Tendinopathy: A

412 Randomized Controlled Trial. *Am. J. Sports Med.* 43, 1704–11. doi:10.1177/0363546515584760

413 Bohm, S., Mersmann, F., Arampatzis, A., 2015. Human tendon adaptation in response to mechanical

414 loading: a systematic review and meta-analysis of exercise intervention studies on healthy adults.

415 *Sport. Med. - Open* 1, 7. doi:10.1186/s40798-015-0009-9

416 Buckley, M., Sarver, J., Freedman, B., Soslowsky, L., 2013. The dynamics of collagen uncrimping

417 and lateral contraction in tendon and the effect of ionic concentration. - PubMed - NCBI. *J*

418 *Biomech* 46, 2242–9.

419 Chimenti, R.L., Flemister, A.S., Tome, J., McMahon, J.M., Flannery, M.A., Xue, Y., Houck, J.R.,

420 2014. Altered tendon characteristics and mechanical properties associated with insertional

421 achilles tendinopathy. *J. Orthop. Sports Phys. Ther.* 44, 680–9. doi:10.2519/jospt.2014.5369

422 Ciccone, W.J., Bratton, D.R., Weinstein, D.M., Elias, J.J., 2006. Viscoelasticity and temperature

423 variations decrease tension and stiffness of hamstring tendon grafts following anterior cruciate

424 ligament reconstruction. *J. Bone Joint Surg. Am.* 88, 1071–8. doi:10.2106/JBJS.E.00576

425 Cook, J.L., Purdam, C.R., 2014. The challenge of managing tendinopathy in competing athletes. *Br. J.*

426 *Sports Med.* 48, 506–9. doi:10.1136/bjsports-2012-092078

427 Drew, B.T., Smith, T.O., Littlewood, C., Sturrock, B., 2012. Do structural changes (eg,

428 collagen/matrix) explain the response to therapeutic exercises in tendinopathy: a systematic

429 review. *Br. J. Sports Med.* 1–8. doi:10.1136/bjsports-2012-091285

430 Dubinskaya, V.A., Eng, L.S., Rebrow, L.B., Bykov, V.A., 2007. Comparative study of the state of

431 water in various human tissues. *Bull. Exp. Biol. Med.* 144, 294–7.

432 Grigg, N.L., Wearing, S.C., Smeathers, J.E., 2012. Achilles tendinopathy has an aberrant strain

433 response to eccentric exercise. *Med. Sci. Sports Exerc.* 44, 12–7.

434 doi:10.1249/MSS.0b013e318227fa8c

435 Grigg, N.L., Wearing, S.C., Smeathers, J.E., 2009. Eccentric calf muscle exercise produces a greater  
436 acute reduction in Achilles tendon thickness than concentric exercise. *Br. J. Sports Med.* 43,  
437 280–283. doi:10.1136/bjism.2008.053165

438 Hannafin, J.A., Arnoczky, S.P., 1994. Effect of cyclic and static tensile loading on water content and  
439 solute diffusion in canine flexor tendons: an in vitro study. *J. Orthop. Res.* 12, 350–6.  
440 doi:10.1002/jor.1100120307

441 Helland, C., Bojsen-Møller, J., Raastad, T., Seynnes, O.R., Moltubakk, M.M., Jakobsen, V., Visnes,  
442 H., Bahr, R., 2013. Mechanical properties of the patellar tendon in elite volleyball players with  
443 and without patellar tendinopathy. *Br. J. Sports Med.* 47, 862–8. doi:10.1136/bjsports-2013-  
444 092275

445 Henninger, H.B., Underwood, C.J., Ateshian, G.A., Weiss, J.A., 2010. Effect of sulfated  
446 glycosaminoglycan digestion on the transverse permeability of medial collateral ligament. *J.*  
447 *Biomech.* 43, 2567–73. doi:10.1016/j.jbiomech.2010.05.012

448 Kang, S., Thordarson, D.B., Charlton, T.P., 2012. Insertional Achilles tendinitis and Haglund’s  
449 deformity. *Foot ankle Int.* 33, 487–91. doi:10.3113/FAI.2012.0487

450 Kongsgaard, M., Qvortrup, K., Larsen, J., Aagaard, P., Doessing, S., Hansen, P., Kjaer, M.,  
451 Magnusson, S.P., 2010. Fibril morphology and tendon mechanical properties in patellar  
452 tendinopathy: effects of heavy slow resistance training. *Am. J. Sports Med.* 38, 749–756.  
453 doi:10.1177/0363546509350915

454 Maffulli, N., Reaper, J., Ewen, S.W.B., Waterston, S.W., Barrass, V., 2006. Chondral metaplasia in  
455 calcific insertional tendinopathy of the Achilles tendon. *Clin. J. Sport Med.* 16, 329–34.

456 Malliaras, P., Cook, J., Purdam, C., Rio, E., 2015. Patellar Tendinopathy: Clinical Diagnosis, Load  
457 Management, and Advice for Challenging Case Presentations. *J. Orthop. Sports Phys. Ther.* 45,  
458 887–98. doi:10.2519/jospt.2015.5987

459 Maquirriain, J., Kokalj, A., 2014. Acute Achilles tendinopathy: effect of pain control on leg stiffness. *J.*  
460 *Musculoskelet. Neuronal Interact.* 14, 131–6.

461 Rio, E., Kidgell, D., Purdam, C., Gaida, J., Moseley, G.L., Pearce, A.J., Cook, J., 2015. Isometric

462 exercise induces analgesia and reduces inhibition in patellar tendinopathy. *Br. J. Sports Med.* 49,  
463 1277–83. doi:10.1136/bjsports-2014-094386

464 Thornton, G., Shrive, N., Frank, C., 2001. Altering ligament water content affects ligament pre-stress  
465 and creep behaviour. - PubMed - NCBI. *J Orthop Res* 19, 845–51.

466 Visentini, P.J., Khan, K.M., Cook, J.L., Kiss, Z.S., Harcourt, P.R., Wark, J.D., 1998. Om ' ns I I I ) 22–  
467 28.

468 Wall, M.E., Banes, A.J., 2005. Early responses to mechanical load in tendon: role for calcium  
469 signaling, gap junctions and intercellular communication. *J. Musculoskelet. Neuronal Interact.* 5,  
470 70–84.

471 Wearing, S.C., Hooper, S.L., Purdam, C., Cook, J., Grigg, N., Locke, S., Smeathers, J.E., 2013. The  
472 acute transverse strain response of the patellar tendon to quadriceps exercise. *Med. Sci. Sports  
473 Exerc.* 45, 772–777. doi:10.1249/MSS.0b013e318279a81a

474 Wearing, S.C., Locke, S., Smeathers, J.E., Hooper, S.L., 2015. Tendinopathy alters cumulative  
475 transverse strain in the patellar tendon after exercise. *Med. Sci. Sports Exerc.* 47, 264–271.  
476 doi:10.1249/MSS.0000000000000417

477 Zwerver, J., Bredeweg, S.W., Hof, a L., 2007. Biomechanical analysis of the single-leg decline squat.  
478 *Br. J. Sports Med.* 41, 264–268; discussion 268. doi:10.1136/bjism.2006.032482  
479